

Original Articles

The prevalence of primary neck and shoulder pain, and its related factors in Japanese postpartum women

K. Koyasu¹, M. Kinkawa², N. Ueyama³, Y. Tanikawa⁴, K. Adachi⁵, H. Matsuo¹

¹ Kobe University Graduate School of Health Sciences, Kobe; ² Ube Frontier University, Ube

³ Takarazuka University School of Nursing, Osaka

⁴ Kansai University of International Studies, Miki; ⁵ Tokyo Metropolitan University, Tokyo (Japan)

Summary

Purpose: This study investigated the prevalence, location, and severity of neck and shoulder pain (NSP), its disturbance of quality of life (QOL), and the factors related to NSP in Japanese postpartum women. **Materials and Methods:** The study involved 308 postpartum women who had a medical examination one month after delivery. The questionnaire consisted of the background and details of NSP. Mood states were evaluated using the Profile of Mood States-Brief (POMS-B), Japanese Version. **Results:** The prevalence of NSP was 73.1%, one-fourth of which occurred after birth. The most common area was the superior part of the trapezius muscles. Prevalence was associated with past history of premenstrual syndrome (PMS), anemia during pregnancy, time per breastfeeding, and the mean POMS-B Fatigue score. Total breastfeeding time a day, the mean POMS-B score for Fatigue, Confusion, Anger-Hostility, and Depression were significantly higher for “worse” after birth than those for “no-change/relief”. The disturbance of daily life due to NSP in postpartum women with past history of PMS and Hiesho were significant higher than that for women without those. **Conclusions:** The prevalence of NSP in postpartum women was very high. The factors which affect NSP were the mental states, breastfeeding, past history of PMS, and anemia during pregnancy.

Key words: Neck and shoulder pain; Postpartum women; Prevalence; Breastfeeding.

Introduction

Neck and shoulder pain (NSP) is the most common symptom for Japanese women in a Japanese comprehensive survey of living conditions conducted by the Ministry of Health, Labour, and Welfare in 2010 [1]. The prevalence of NSP in Japanese women in their 30s is 14.6%, which is double than that for men, and it rises with age. NSP is often accompanied by a pain and an unpleasant symptom and the pain reduces quality of life (QOL) [2].

NSP is classified into primary and secondary pain. Primary NSP is defined as the absence of a definite disease, while secondary NSP defined as the presence of a definite disease.

It has been demonstrated that primary NSP may be associated with multiple factors, including smoking, obesity [3], women, age [4, 5], working conditions [4, 6-8], and psychological distress such as depression or anxiety [9-13].

Postpartum women constantly play many important roles and are in a physically and mentally stressful condition [14, 15], which may be related to the prevalence or severity of NSP. However, there are no reports demonstrating what kind of factors affect the prevalence and severity of NSP, and its QOL disturbance in postpartum women.

Thus, the authors conducted the present study to examine the prevalence, location, and severity of NSP and its disturbance of QOL in Japanese postpartum women, and to assess the kind of factors related to NSP.

Materials and Methods

Subjects

This study was conducted at two hospitals and one obstetric clinic in Kobe city, during October 2011 and April 2012.

Subjects were postpartum women with both term births and normal newborn babies. The women were excluded if they had an orthopedic disease.

Questionnaires were distributed to the postpartum women who consented to this study at their medical examination one month after delivery, and these were then deposited into a special box beside the reception desk (collection rate: 83.7%).

This study was approved by the Ethical Committee at Kobe University Graduate School of Health Sciences.

Self-administered questionnaire

The questionnaire consisted of the subject's background and details of the NSP. Background included age, height, weight, delivery history, delivery style, period after birth, anemia during pregnancy and after delivery, past history of premenstrual syn-

Table 1. — *Subject characteristics (n = 308).*

Background	Mean \pm SD or n (%)	Range
Age (years)	31.9 \pm 5.1	18-43
BMI (kg/m ²)	20.4 \pm 2.7	
Day after birth	35.5 \pm 7.3	17-58
Delivery history		
primipara	148 (48.1)	
multipara	160 (51.9)	
Delivery style		
vaginal delivery	254 (82.5)	
caesarean section	54 (17.5)	
Anemia during pregnancy	146 (47.4)	
Anemia after birth	87 (28.2)	
History of PMS	63 (20.5)	
Hieshou	199 (64.6)	
Breastfeeding method		
breastfeeding only	237 (76.9)	
breastfeeding and bottle-feeding	64 (20.8)	
bottle-feeding only	7 (2.3)	

BMI: body mass index; PMS: premenstrual syndrome

drome(PMS) and Hieshou, and method, position, frequency, and duration of breastfeeding.

NSP included the present history, onset (before and during pregnancy, after birth), change of NSP after birth (five levels: worse, slightly worse, no change, a little relief, relief), areas for NSP, daily living activities which made the NSP worse, and disturbance of daily life due to NSP (level 0 (none) to 10).

Profile of Mood States-Brief (POMS-B), Japanese version

Mood states were evaluated using the 30-item POMS-B, Japanese version. POMS-B is able to measure temporary change in feeling according to condition. The subject answered for each item her mood over the past one week.

The answer to each question was described using a five-point scale as: not at all, a little, moderately, quite a bit or extremely. A

score of 0 to 4 was then assigned to each answer. POMS-B consists of the six mood state: “Tension-Anxiety”(T-A), “Depression”(D), “Anger-Hostility”(A-H), “Fatigue”(F), “Vigor”(V), and “Confusion”(C). The scores were the sums of the items for each mood state and these were calculated as a T-score. T-score = $50 + 10 \times (\text{score} - \text{mean}) / \text{SD}$.

The scores of the aforementioned items were compared between two groups: with NSP and without NSP, and worse after birth (worse, slightly worse) and no change/relief after birth (no change, a little relief, relief), for NSP, respectively.

Statistics

The differences in background, present NSP, and POMS-B score between the two groups were tested using the t-test and χ^2 test. Statistical significance was expressed as *p* values at 95% confidence intervals. All statistical analyses were carried out using SPSS for Windows (20J).

Results

Characteristics of subjects

Table 1 shows the subject characteristics. The mean of age was 31.9 ± 5.1 years (range 18–43). The mean of BMI was 20.4 ± 2.7 kg/m². The percentage of primipara and multipara were 48.1% (148/308) and 51.9% (160/308).

The rates of anemia during pregnancy and after birth were 47.4%(146/308) and 28.2% (87/308). Postpartum women with a past history of PMS were 20.5% (63/308) and those with a Hieshou were 64.6% (199/308). Concerning breastfeeding, breastfeeding only, breastfeeding and bottle-feeding, and bottle-feeding only, were 76.9% (237/308), 20.8% (64/308), and 2.3% (7/308), respectively.

NSP in postpartum women

1) Prevalence of NSP:

A total of 225 individuals answered “yes” to the question “Do you presently have NSP?” The prevalence of NSP was 73.1% (225/308)




Figure 1. — Present history of neck and shoulder pain. A) Onset of neck and shoulder pain in postpartum women. B) Change of neck and shoulder pain after birth in postpartum women.




Figure 2. — The areas for NSP. A) Distribution of areas for NSP. B) The strongest area for NSP.

2) Present history of NSP (onset, change after birth):

Onset of NSP was categorized into four categories: before pregnancy, during pregnancy, after birth, and others, and these were 66.2% (149/225), 4.0% (9/225), 24.9% (56/225), and 4.9% (11/225), respectively (Figure 1A). The mean of onset of NSP after birth was 8.1 ± 5.9 days. Change of NSP after birth: “worse” and “slightly worse” were about 44.5% (100/225), “no change” was reported by 40.0% (90/225), “a little relief” by 12.4% (28/225), and “relief” by 3.1% (7/225) (Figure 1B).

3) The areas of NSP (Figure 2):

The most common areas were the superior part of the trapezium muscles. The strongest area was the left posterior cervical region 25.8% (58/225), and the next one was the left superior part of trapezium 25.3% (57/225) (Figure 2B).




Figure 3. — Daily living activities that cause NSP to become worse.

4) Daily living activities which made NSP worse:

A total of 92.9% (209/225) of postpartum women with NSP reported that it became worse through daily living activities. The most frequently reported daily living activities were breastfeeding 69.3% (156/225), holding the baby 60.4% (136/225), and using a computer 37.8% (85/225) (multiple answer) (Figure 3).

Factors affecting NSP

1) Factors related to prevalence of NSP:

The prevalence of NSP was associated with past history of PMS ($p = 0.000$) and anemia during pregnancy ($p = 0.032$). Hiesho, anemia after delivery, smoking, age, delivery history, and delivery style did not appear to have a significant influence on NSP. There was a significant difference in the mean duration per breastfeeding (minutes) between the two groups ($p = 0.041$). However method and position did not appear to have a significant influence on the prevalence of NSP.

Regarding the T-scores of POMS-B, the mean Fatigue (F) score for postpartum women with NSP was significantly higher than for those without NSP ($p = 0.003$). The postpartum women with NSP showed higher scores in Tension-Anxiety (T-A), Depression (D), Anger-Hostility (A-H), and Confusion (C), and a lower score for Vigor (V), compared with those without neck and shoulder pain, but there were no significant differences (Table 2).

2) Factors related to change of NSP after birth:

There were no significant differences in anemia during pregnancy, anemia after delivery, past history of PMS, Hiesho, and delivery style between the two groups. In regards to breastfeeding, total time per day was significantly longer in the “worse” group than the “no change/relief” group ($p = 0.018$). The method, position, frequency per day, and breast tension did not appear to have a significant influence on causing the NSP to become worse after birth.

Table 2. — *Present NSP (n = 308).*

	Mean \pm SD or n		
	yes (225)	no (83)	<i>p</i>
Age (years)	31.9 \pm 5.0	31.8 \pm 5.2	0.805
BMI (kg/cm ²)	20.4 \pm 2.6	20.6 \pm 3.0	0.504
Delivery history			
primipara	115	33	0.077
multipara	110	50	
Anemia during pregnancy			
yes	115	31	0.032*
no	110	52	
Anemia after birth			
yes	69	18	0.120
no	156	65	
Delivery style			
vaginal delivery	183	71	0.389
caesarean section	42	12	
History of PMS			
yes	57	6	0.000**
no	168	77	
Hiesho			
yes	151	48	0.131
no	74	35	
Smoking			
non-smoker	197	73	0.925
current/ex-smoker	28	10	
Breastfeeding method			
breastfeeding only	167	70	0.061
breastfeeding and bottle-feeding / bottle-feeding	58	13	
only			
Breastfeeding			
frequency (times / day)	9.6 \pm 2.1	9.6 \pm 2.3	0.968
one time (minutes)	20.9 \pm 10.5	18.2 \pm 7.9	0.041*
total times a day (minutes)	198.5 \pm 106.0	176.6 \pm 93.5	0.113
POM-B			
T-A	44.4 \pm 8.7	42.8 \pm 8.3	0.146
D	44.7 \pm 7.2	44.2 \pm 6.8	0.645
A-H	48.4 \pm 8.9	44.7 \pm 9.8	0.564
V	44.7 \pm 9.7	46.7 \pm 9.4	0.094
F	47.8 \pm 7.8	44.8 \pm 8.3	0.003**
C	43.1 \pm 7.9	43.2 \pm 8.1	0.919

* Correlation is significant at the 0.05 level;

**Correlation is significant at the 0.01 level.

Table 3. — *Change of NSP after birth (n = 225).*

	Mean \pm SD or n		
	worse (100)	no-change / relief (125)	<i>p</i>
Age (years)	32.0 \pm 5.1	31.9 \pm 5.0	0.920
BMI (kg/cm ²)	20.1 \pm 2.4	20.5 \pm 2.8	0.504
Delivery history			
primipara	53	62	0.612
multipara	47	63	
Anemia during pregnancy			
yes	44	71	0.056
no	56	54	
Anemia after birth			
yes	35	34	0.207
no	65	91	
Delivery style			
vaginal delivery	87	96	0.051
caesarean section	13	29	
History of PMS			
yes	29	28	0.258
no	71	97	
Hiesho			
yes	67	84	0.975
no	33	41	
Smoking			
non-smoker	87	110	0.821
current/ex-smoker	13	15	
Breastfeeding method			
breastfeeding only	71	96	0.323
breastfeeding and bottle-feeding / bottle-feeding	29	29	
only			
Breastfeeding			
frequency (times / day)	9.8 \pm 2.2	9.4 \pm 2.1	0.238
one time (minutes)	22.4 \pm 11.7	19.6 \pm 9.2	0.058
total times a day (minutes)	218.0 \pm 116.7	182.3 \pm 93.6	0.018*
POM-B			
T-A	44.7 \pm 9.6	44.1 \pm 8.0	0.585
D	45.9 \pm 8.9	43.6 \pm 5.4	0.023*
A-H	49.9 \pm 10.0	47.2 \pm 7.8	0.030*
V	44.3 \pm 9.6	44.9 \pm 9.8	0.567
F	49.9 \pm 8.5	46.2 \pm 6.8	0.000**
C	45.0 \pm 9.3	41.5 \pm 6.1	0.002**

* Correlation is significant at the 0.05 level;

**Correlation is significant at the 0.01 level.

Regarding the T-scores of POMS-B, the mean score for D, A-H, F, and C were significantly higher for “worse” than for “no-change/relief” of NSP ($p = 0.023$, $p = 0.030$, $p = 0.000$, and $p = 0.002$, respectively). This study showed that the T-A and V scores of postpartum women who perceived worse pain did not differ significantly from those who perceived no-change/relief after birth (Table 3).

Disturbance of daily life due to NSP and related factors

The mean score for postpartum women with NSP was 4.6 \pm 2.3 (Figure 4). Table 4 shows the disturbance of daily life due to NSP, and its related factors. The mean scores for

“worse” were significantly higher than for “no change/relief” after birth ($p = 0.001$). There were significant differences in the mean scores for disturbance of daily life between Hiesho and past history of PMS ($p = 0.043$, $p = 0.020$).

Discussion

This is the first instance to demonstrate that the prevalence of primary NSP in postpartum women was 73.1%, one-fourth of which was after birth-onset, and that the most common area was the superior part of the trapezium muscles, while the strongest area was the left posterior cervical re-




Figure 4. — Disturbance of daily life due to NSP in postpartum women.

gion. The factors which affect NSP in postpartum women were mental states, breastfeeding, anemia during pregnancy and past history of PMS. Breastfeeding was the most common daily living activity that caused NSP to become worse.

The present research demonstrated that the prevalence of NSP in Japanese postpartum women was 73.1%. In Sweden, it was found to be 29.4% in four to eight weeks after childbirth [16]. Hill *et al.* [3] reported that 22.3% of participants who were over 18 years of age had pain, aching or stiffness in either shoulders. Hakala *et al.* [5] demonstrated that pain of the neck and shoulder affected 45% in 18-year-old girls. Therefore, the prevalence of NSP for Japanese postpartum women could be higher than that for postpartum women of other countries or those of other generations.

The most common areas for NSP in postpartum women in our study were the superior part of the trapezium muscles and the posterior region of the neck. This finding coincided with that for nurses reported by Iizuka *et al.* [11]. However, the rate of NSP in the posterior cervical region for postpartum women was higher than that for nurses. The muscle of the posterior cervical regions were extended by anteflexion posture during breastfeeding. This posture might cause NSP. Actually, the rate of after birth-onset NSP in postpartum women had a high prevalence in our study. Thus, it was suggested that breastfeeding might contribute to the increase in NSP after birth-onset.

Interestingly, NSP in postpartum women appeared outstandingly stronger on the left side compared with the right side. Fujii *et al.* [17] reported that the strongest areas for neck-shoulder discomfort in employees were the right upper scapula 44.4% and the right neck 29.6%. It was noted that postpartum women put the head of the baby on the side opposite their handedness (i.e. the left side for right handed women) for breastfeeding. It was therefore speculated that breastfeeding caused the higher rate for the left side NSP in postpartum women.

In our study, the mean POMS-B F score for postpartum women with NSP was significantly higher than that for those without NSP. The mean score for D, A-H, F and C were sig-

Table 4. — Disturbance of daily life due to NSP (n=225).

	Mean±SD	P
Change of neck and shoulder pain after birth		
worse	5.1±2.4	
no-change/relief	4.1±2.1	0.001**
Heisho		
yes	4.8±2.2	
no	4.1±2.4	0.043*
History of PMS		
yes	5.2±2.2	
no	4.4±2.3	0.020*

* Correlation is significant at the 0.05 level;

**Correlation is significant at the 0.01 level.

nificantly higher for “worse” than that for “no-change/relief” after birth. Several research studies have reported that NSP is related to psychological factors such as depression [10,18], confusion [19], working stress and the extent of the feeling of satisfactions [8,13]. Psychological stress such as anxiety or depression in postpartum women may be more than that for others [20]. It was suggested that psychological states could affect the onset of NSP and cause it to worsen after birth in postpartum women.

In this study, about 70% of postpartum women with NSP answered that breastfeeding made it worse in their daily life activities. In addition, breastfeeding was carried out nine times a day, taking 20 minutes each time, for a total of 190 minutes a day. In particular, the required time per day for breastfeeding was related to the worsening of NSP after birth. Muscles around the neck and the shoulder girdle support the head and the arms, which weigh more than four kg, therefore there is always a load on the muscle around the shoulder girdle from the neck [21]. The posture of postpartum women tends to become unbalanced because of not-held head of the babies, and due to keeping the anteflexion posture for feeding. It seems likely that the after birth worsening of NSP may be related to muscles strain, due to the posture adopted for breastfeeding and the length of time it takes.

There was significant difference in QOL level in the disturbance of daily life for “worse” and “no-change/relief” NSP in postpartum women. The number of involved sites of self-reported musculoskeletal pain was associated with the level of reduction in health-related quality of life (HRQOL) among young adults [2]. Respondents with shoulder pain scored lower on all domains of the SF36 [3]. Thus, NSP after birth led to the reduction in the level of QOL. The disturbance of daily life due to NSP in postpartum women with past history of PMS and Hieshou was significantly higher than those without those. PMS and Hieshou are health of obstacles related to the ovarians steroid hormones cyclist [22,23]. We speculated that the sudden hormone change after birth might contribute to the physiology of NSP.

Conclusions

This study demonstrated that the prevalence of NSP in postpartum women was very high, and the rate of after birth-onset and worsening after birth were very high. In addition, NSP after birth led to a reduction in the level of QOL of postpartum women. It was revealed that NSP greatly influenced the health of postpartum women. The factors which affect NSP were related to the mental states of postpartum women, time per breastfeeding, anemia during pregnancy, and past history of PMS. Breastfeeding in particular was identified by postpartum women as the daily living activity most responsible for causing their NSP to become worse. It was therefore suggested that this was an important factor.

Further studies should involve the collection and analysis of objective data to clarify how these factors influence the mechanism of NSP. It is also necessary to examine care for the prevention and improvement of NSP in postpartum women.

Acknowledgments

The authors are grateful to all of the postpartum women who took part in this study and all those concerned who supported this study.

References

- [1] Statistics and Information Department, Ministry of Health, Labour and Welfare: "Comprehensive survey of living conditions of the people on health and welfare 2010. Ministry of Health, Labour and Welfare". Available at: <http://www.mhlw.go.jp/toukei/saikin/hw/k-tyosa/k-tyosa10/toukei.html>. Accessed on 2011.10.1.
- [2] Paananen M., Taimela S., Auvinen J., Tammelin T., Zitting P., Karpinen J.: "Impact of self-reported musculoskeletal pain on health-related quality of life among young adults". *Pain Med.*, 2011, 12, 9.
- [3] Hill C.L., Gill T. K., Shanahan E. M., Taylor A.W.: "Prevalence and correlates of shoulder pain and stiffness in a population-based study: the North West Adelaide Health Study". *Int. J. Rheum. Dis.*, 2010, 13, 215.
- [4] Cassou B., Derriennic F., Monfort C., Norton J., Touranchet A.: "Chronic neck and shoulder pain, age, and working conditions: longitudinal results from a large random sample in France". *Occup. Environ. Med.*, 2002, 59, 537.
- [5] Hakala P., Rimpela A., Salminen J.J., Virtanen S.M., Rimpela M.: "Back, neck, and shoulder pain in Finnish adolescents: national cross sectional surveys". *BMJ.*, 2002, 325, 743.
- [6] Ariens G.A., Bongers P.M., Douwes M., Miedema M.C., Hoogendoorn W.E., van der Wal G.: "Are neck flexion, neck rotation, and sitting at work risk factors for neck pain? Results of a prospective cohort study". *Occup. Environ. Med.*, 2001, 58, 200.
- [7] Chee H.L., Rampal K.G., Chandrasakaran A.: "Ergonomic risk factors of work processes in the semiconductor industry in Peninsular Malaysia". *Ind. Health*, 2004, 42, 373.
- [8] Leino P., Magni G.: "Depressive and distress symptoms as symptom as predictors of low-back-pain, neck-shoulder-pain, and other musculoskeletal morbidity: A 10-year follow-up of metal-industry employees". *Pain*, 1993, 53, 89.
- [9] Diepenmaat A.C. M., van der Wal M.F., de Vet H.C.W., Hirasings R.A.: "Neck/shoulder, low back, and arm pain in relation to computer use, physical activity, stress, and depression among Dutch adolescents". *Pediatrics*, 2006, 117, 412.
- [10] Ishizuka B., Kudo Y., Tango T.: "Cross-sectional community survey of menopause symptoms among Japanese women". *Maturitas*, 2008, 61, 260.
- [11] Iizuka Y., Shinozaki T., Kobayashi T., Tsutsumi S., Osawa T., Ara T.: "Characteristics of neck and shoulder pain (called katakori in Japanese) among members of the nursing staff". *J. Orthop. Sci.*, 2012, 17, 46.
- [12] Linton S.J., Kamwendo K.: "Risk-factors in the psychosocial work-environment for neck and shoulder pain in secretaries". *J. Occup. Environ. Med.*, 1989, 31, 609.
- [13] Holmstrom E.B., Lindell J., Moritz U.: "Low-back and neck shoulder pain in construction workers-occupational workload and psychosocial risk-factors. 2.Relationship to neck and shoulder pain". *Spine*, 1992, 17, 672.
- [14] Dennis C.L., Ross L.: "Relationships among infant sleep patterns, maternal fatigue, and development of depressive symptomatology". *Birth*, 2005, 32, 187.
- [15] Thompson J.F., Roberts C.L., Currie M., Ellwood D.A.: "Prevalence and persistence of health problems after childbirth: Associations with parity and method of birth". *Birth*, 2002, 29, 2, 83.
- [16] Schytt E., Lindmark G., Waldenstrom U.: "Physical symptoms after childbirth: prevalence and associations with self-rated health". *BJOG*, 2005, 112, 210.
- [17] Fujii T., Matsudaira K., Noma K., Ishizuka A., Yamada K., Arisaka M.: "Objective measurement of neck-shoulder discomfort and analysis of associated factors". *Clin. Orthop. Surg.*, 2012, 47, 619.
- [18] Vogt M.T., Simonsick E.M., Harris T.B., Nevitt M.C., Kang J.D., Rubin S.M.: "Neck and shoulder pain in 70-to 79-year-old men and women: findings from the health, aging and body composition study". *Spine J.*, 2003, 3, 435.
- [19] Makela M., Heliovaara M., Sievers K., Impivaara O., Knekt P., Aromaa A.: "Prevalence, determinants, and consequences, of chronic neck pain in Finland". *Am. J. Epidemiol.*, 1991, 134, 1356.
- [20] Josefsson A., Berg G., Nordin C., Sydsjo G.: "Prevalence of depressive symptoms in late pregnancy and postpartum. *Acta Obstet. Gynecol. Scand.*, 2001, 80, 251.
- [21] Takagishi K., Hoshino Y., Ide J., Sugihara T., Hata Y., Sano H.: "Project study on katakori (2004-2006)". *J. Jpn. Orthop. Assoc.*, 2008, 82, 901.
- [22] Mortola J.F.: "Premenstrual syndrome". *Trends Endocrinol Metab*, 1996, 7, 184.
- [23] Orimo A., Inoue S., Ikegami A., Hosoi T., Akishita M., Ouchi Y.: "Vascular smooth-muscle cells as target for estrogen". *Biochem. Biophys. Res. Commun.*, 1993, 195, 2730.

Address reprint requests to:

H. MATSUO, M.D.

Kobe University Graduate School of Health Sciences

7-10-2 Tomogaoka, Suma-ku,

Kobe 654-0142 (Japan)

e-mail: matsuo@tiger.kobe-u.ac.jp