

Acquired prosopagnosia: structural basis and processing impairments

Jodie Davies-Thompson¹, Raika Pancaroglu¹, Jason Barton¹

¹*Department of Medicine (Neurology), and Department of Ophthalmology and Visual Sciences, University of British Columbia, Canada*

TABLE OF CONTENTS

1. Abstract
2. Introduction
3. Acquired prosopagnosia
 - 3.1. Apperceptive prosopagnosia
 - 3.2. Associative/amnestic prosopagnosia
 - 3.3. Impairments of other functions
 - 3.4. Face specificity versus expertise
 - 3.5. Covert face recognition
4. Developmental prosopagnosia
5. Summary
6. Acknowledgments
7. References

1. ABSTRACT

Cognitive models propose a hierarchy of parallel processing stages in face perception, and functional neuroimaging shows a network of regions involved in face processing. Reflecting this, acquired prosopagnosia is not a single entity but a family of disorders with different anatomic lesions and different functional deficits. One classic distinction is between an apperceptive variant, in which there is impaired perception of facial structure, and an associative/amnestic variant, in which perception is relatively intact, with subsequent problems matching perception to facial memories, because of either disconnection or loss of those memories. These disorders also have to be distinguished from people-specific amnesia, a multimodal impairment, and prosop-anomia, in which familiarity with faces is preserved but access to names is disrupted. These different disorders can be conceived as specific deficits at different processing stages in cognitive models, and suggests that these functional stages may have distinct neuroanatomic substrates. It remains to be seen whether a similar anatomic and functional variability is present in developmental prosopagnosia.

2. INTRODUCTION

Prosopagnosia is the impaired ability to recognize familiar faces or to learn to recognize new faces (1, 2). Individuals with prosopagnosia have a problem discriminating known from unknown faces, and in the absence of useful information about face identity they experience most faces as unfamiliar (which contrasts with the bias to experiencing most faces as familiar that is seen in patients with “false familiarity for faces” (3). Because of the pre-eminence of faces in social interactions, prosopagnosia is one of the most well-known and well-studied deficits of person recognition. Although impaired face recognition can be a symptom of more general problems of perception, cognition, and memory, as in macular degeneration (4), Alzheimer’s dementia (5-7), mild cognitive impairment (8), Huntington’s chorea (9), and Parkinson’s disease (10, 11), the term ‘prosopagnosia’ should be reserved for a specific dysfunctional state in which there is a relatively selective deficit in face recognition that cannot be explained by more general problems of perception, cognition or memory. Therefore, diagnosing this condition requires detailed

Acquired prosopagnosia: structure and perception




Figure 1. Model of face processing. Damage to posterior regions of the inferior occipito-temporal cortex can result in difficulties with early perceptual encoding and perceptual encoding of static structure, resulting in apperceptive prosopagnosia (blue). Damage to the anterior inferior temporal cortex can result in difficulties retrieving face memories, or linking static structure to the memory of faces (red). Meanwhile, damage to the anterior temporal pole can result in difficulties with biographical information which results in person-specific amnesia (green).

neuropsychological evaluation of these general functions as well as tests that assess either long-term recognition of familiar faces, as with a variety of familiar faces tests that use images of celebrities (12), or short-term memory formation for new faces, as with the Warrington Recognition Memory Test (13) or the Cambridge Face Memory Test (14).

Why and how people can acquire a deficit that seems to affect mainly the ability to recognize faces is the subject of substantial research. One important point deserving emphasis is that prosopagnosia is not a disorder but a family of disorders. Face recognition is a complex cognitive process, and like all such processes it cannot be reduced to a single function or a single anatomic region in the brain. Rather, it requires participation of several perceptual, memory and cognitive operations that take place in a distributed network of interconnected cerebral regions. As a result, the lesions of acquired prosopagnosia can vary significantly between patients, as can the precise nature of their dysfunction and associated defects.

Structurally, it has long been known that prosopagnosia can arise from a very diverse set of lesions, ranging in size from posterior occipitotemporal regions to anterior temporal cortex (15). While early reports described prosopagnosia after bilateral occipitotemporal lesions (16, 17), others suggested that a right hemispheric lesion could be sufficient to cause prosopagnosia (18, 19). This was supported by later work with neuroimaging indicating that some patients had unilateral lesions, almost always on the right side (19, 20). Of the few subjects with unilateral left lesions, most were left-handed (21-24), suggesting that they may have had anomalous hemispheric asymmetries to

begin with. A right hemispheric bias for cases of prosopagnosia accords with data from functional magnetic resonance imaging studies in healthy subjects, which show that faces activate more voxels with greater significance in the core face-processing network of the right hemisphere, compared to those components in the left hemisphere (25, 26). At this point, though, it is not clear how prosopagnosia with bilateral lesions differs from prosopagnosia with unilateral right-sided lesions. Although there is some suggestion from functional neuro-imaging that the components of the face network in the left hemisphere may be involved more with perception of local features of the face (27), it is not known for certain whether these components perform redundant operations or forms of processing that are complementary to their right-sided counterparts. Nevertheless, there are occasional reports of patients who suffered right-side damage but did not report prosopagnosia until after a second left-sided lesion (28), suggesting that in at least some human subjects the contribution of the left hemisphere to face recognition is significant.

Functionally, concepts of the nature of the dysfunction in prosopagnosia are closely linked to models of normal face recognition. One enduring and influential model (29) contains a number of key aspects. First, a prominent feature is the proposal of two parallel mechanisms – one involved in processing the changeable aspects of faces, such as emotional expression, viewpoint, lip-reading, and gaze direction, and another involved in the perception of stable aspects of faces, particularly identity (Figure 1). This is supported by evidence from functional imaging and primate electrophysiology that there may indeed be separate areas for encoding facial identity and

Acquired prosopagnosia: structure and perception

facial social signals (30-32). If so, this raises the possibility that deficits in the processing of identity and social signals could be dissociated in patients with lesions, and that not all types of facial information may be affected in prosopagnosia. Although the literature is mixed on whether prosopagnosic patients are also impaired on aspects of face perception like expression and lip-reading (see (33) for a review), there is some evidence for a double dissociation between identity and expression processing (34).

A second important feature is that face processing involves a hierarchy of stages (35, 36). In the case of identity, visual processing generates a face percept, which is then matched to a memory store of previously encountered faces, sometimes referred to as 'face recognition units', or FRUs. A successful match activates person-identity nodes (PINs), which in turn activate name recognition units (NRUs) and semantic information units (SRUs) containing biographical information about the person. PINs, SRUs and NRUs are multimodal in nature, in that they can also be accessed through other non-facial cues, such as voice or gait. Top-down processing can also occur. For example, matching of the face percept to face recognition units may be enhanced by top-down semantic activation from person identity nodes, as when the subject already knows the name of the person or some contextual information about them (e.g. actor, family, workmate).

Reflections of this hierarchical model are also evident in some of the clinical aspects of prosopagnosia. First, there is evidence of functional subtypes of prosopagnosia, in which either the creation of the face percept or its subsequent matching to facial memories is the critical deficit (15, 37, 38). Second, patients with prosopagnosia can identify people by voices or non-facial visual cues such as gait or mannerisms, indicating preserved access to names and semantic information about people through routes other than face perception. In fact, this ability should be part of the diagnostic criteria for the condition, to differentiate prosopagnosia from a multi-modal 'people-specific amnesia' (39, 40). Third, they often describe effects related to top-down processing, in that they can sometimes recognize faces when contextual cues narrow the range of possibilities, such as knowing which people will be at a meeting or are likely to be encountered in a specific setting (41-43).

One of the important goals of research in prosopagnosia is to correlate the structural variations with functional subtypes, to better understand the anatomic basis of the different cognitive processes involved. Most of the evidence that has been collected has been performed on cases with acquired prosopagnosia, which will be the major focus of this article. A developmental form of prosopagnosia has only been recognized more recently: it is increasingly a topic of study, and is more extensively reviewed in the article of Avidan and Behrmann in this issue.

3. ACQUIRED PROSOPAGNOSIA

Acquired prosopagnosia is the loss of the ability to recognize familiar faces following some cerebral injury (44). Most often this affects both the recognition of familiar

faces as well as new faces. However, there is a rare anterograde form, in which the deficit is limited to faces encountered since the onset of the lesion (45-47).

As mentioned above, whether or not the face-processing deficit is limited to the recognition of identity is of theoretical interest to cognitive models. The older literature is replete with contradictory claims. Some prosopagnosic subjects are purportedly impaired in processing gaze direction, emotional expression, age, ethnicity and gender (41, 42, 48-51), while in others the face processing defect is stated to spare at least some of these other types of facial information (39, 52-55). One simple explanation for these divergent claims may be variations in lesion extent and location: however, assessing this is difficult with the information provided in older cases from different researchers. Recent advances in neuroimaging and the ability to identify the functional network involved in face-processing in single subjects (56) will advance this issue. A report that used functional magnetic resonance imaging to describe the impact of lesions on the face network noted a double dissociation (Figure 2). This tested the ability of subjects to discriminate changes in faces morphed along either an identity or expression continuum, with the level of difficulty kept equivalent between the two. The updated data from this study shows that, while a few subjects are impaired in neither or impaired in both, some patients do show dissociated performance on this test. Prosopagnosic patients can be impaired in discriminating changes in identity but not changes in expression in morphed facial images. Conversely, a patient with a lesion that eliminated the right superior temporal sulcus had the reverse deficit (34).

The existence of at least two functional subtypes of prosopagnosia is best established for this acquired form (15, 37, 38). In '*apperceptive prosopagnosia*' the defect is an inability to form a sufficiently accurate representation of the face's structure from visual data. This results in a degraded match to facial memory stores, with weak activation that may not be sufficient to achieve a threshold to trigger either a familiarity signal or access to further information about the person. In contrast, in '*associative prosopagnosia*' the patient can perceive facial structure well; rather, the defect lies in the process of matching this high-fidelity facial information to facial memories. In some cases it is postulated that this is due to a disconnection between perception and memory (57). In other cases, it may be simply because facial memories have been lost, a scenario which some would term an '*amnestic variant*' of prosopagnosia (15). Of course, given the variable location and extent of naturally occurring lesions in humans, and the fact that complex cognitive operations are seldom reducible to single anatomic loci, pure versions of these subtypes may be the exception rather than the rule. Indeed, the data show that patients with severe perceptual deficits in face perception often have milder impairments in face imagery, while those with severe impairments of memory for faces have milder problems on perceptual discrimination tests for faces (38). This is consistent with the concept that face recognition emerges from the coordinated action of a




Figure 2. Contrast between recognition of identity and expression in morphed facial images. Used with permission from (34). The four prosopagnosic patients in the yellow zone have impaired perception of morphing-induced changes in identity, but are accurate in discriminating similar changes in expression. In the blue zone is one subject with a lesion of the right superior temporal sulcus, who had the reverse, impaired perception of expression changes but intact discrimination of identity changes. A number of prosopagnosic subjects are normal on both tests (clear area), including some patients with occipitotemporal lesions. (However, some of these admitted later to using a feature-based strategy to solve the identity part of the test, R-IOT4 by concentrating on the eyebrows, for example.) Only two prosopagnosic subjects had problems with both identity and expression (pink zone). L-IOT1 had a left fusiform resection for epilepsy but also had an atrophic right fusiform region: he showed activation by faces of only the left occipital face area and the right superior temporal sulcus of his core network. B-ATOT1 had extensive bilateral fusiform and anterior temporal lesions from herpes encephalitis.

distributed network, though one in which the different nodes of the network have different contributions. Thus, the terms *apperceptive* and *associative/amnestic* may be best used to indicate the predominant rather than the sole functional deficit in a given patient, with an anterior-posterior gradient in which occipital lesions cause more problems with perception than memory for faces, while anterior temporal lesions cause more problems with memory than with perception of faces, as discussed below.

3.1. Apperceptive prosopagnosia

In *apperceptive prosopagnosia*, the subject cannot form an accurate perceptual representation of the facial structure that is specific to identity. This can occur following either unilateral (most often right) or bilateral occipitotemporal damage (Figure 3). Older reports suggest that this is most often correlated with damage to the right or bilateral fusiform gyri (37, 58). More recent evidence from structural MRI suggests that the responsible lesions often occur in the vicinity of the right fusiform face area (59, 60), an assertion that receives support from functional MRI studies confirming loss of the right fusiform face area in some patients (34). However, others have suggested that the most common lesioned site for prosopagnosia is around the right OFA (61), and is supported by patient findings such as PS, whose lesions involved the right occipital face area and the left fusiform face area, but spared the right fusiform face area (62, 63). This is consistent with face recognition emerging from activity across a cortical network rather than a single region, and suggests that both

the occipital and fusiform face areas contribute to the perceptual encoding of faces.

Apperceptive prosopagnosia is diagnosed by showing that patients are impaired not only in familiarity or recognition of known faces, but also impaired in perceiving the difference between faces. The Benton Face Recognition Test (64) asks a subject to choose which face in an array of six images matches a target face, in some items across changes in viewpoint and lighting. However, this test can be failed by non-prosopagnosic patients (65-67), and some prosopagnosic patients may achieve normal accuracy rates but take a long time to do the test, suggesting abnormal perceptual strategies (68). The Cambridge Face Perception Test (69) presents a face at the top of the screen and asks a subject to arrange in order of resemblance six faces that a morphing program has made gradually more different from the top face, within a time limit. Experimental tests can also examine the amount of difference between morphed faces required to support discrimination (34), or examine the ability of subjects to perceive changes to facial shape (70).

The precise nature of the processing deficit that can result in impaired perception of faces but leave basic object recognition intact remains unclear. One hypothesis is that some sort of holistic perceptual mechanism is disrupted. A number of observations in healthy subjects show substantial influences of whole-object structure on face perception (71-74) and studies have shown that some prosopagnosic patients are deficient in the perception of




Figure 3. Axial images from T1-weighted magnetic resonance imaging of two patients with ‘apperceptive’ prosopagnosia. A) Unilateral inferior occipital damage in a 61 year-old man (R-IOT4) presenting with prosopagnosia, topographagnosia, and left quadrantanopia. B) MR axial images of bilateral inferior occipital damage in a 60 year-old man (B-IOT3) presenting with prosopagnosia, topographagnosia, dyschromatopsia and right hemianopia.

whole faces and instead rely on a feature-by-feature strategy in face perception (58, 75-77). Another related hypothesis is that the configuration or spatial relations between individual facial features in this whole-object structure may be particularly important in specifying the structure of a unique face (78-80). In monkeys, electrophysiological recordings have shown that cells in inferior temporal cortex are selective for faces that differ in these spatial relations (81, 82). Studies have shown that prosopagnosic patients with occipitotemporal lesions are impaired in perceiving the configuration of facial features (38, 83, 84). Ultimately, though, it may be that these spatial relations are merely a convenient metric for studying the precision of a subject’s perception of the complex shape of faces, which need to be processed rapidly and in aggregate across the entire face to support efficient recognition of identity.

A third aspect of face perception is the fact that some regions are more salient than others (85). Numerous studies show that subjects pay more attention to the eyes than to the lower face, particularly when the subject is trying to identify the face (86), and that the eye region contains the information most useful for identification (87-91). Prosopagnosic patients may be particularly impaired in using information from the eye region. Their scanning eye movements show a reduced number of fixations in the eye region (92), their discrimination of structural changes in the eye region is particularly poor (93), and they are more

impaired in deducing identity from the eyes than from the mouth (60, 94-96). Loss of normal patterns of saliency in how prosopagnosic subjects process faces may reflect loss of optimal perceptual strategies in deriving identity, which normally should emphasize the eyes.

Whether deficits in perception of more elementary aspects of vision may contribute to prosopagnosia has been considered. Brightness and contrast are important cues to the orientation of surfaces when lighting casts shadows, and therefore are potent sources of information about the shape of complex objects with multiple surfaces. Studies with photonegatives confirm an important role for brightness information in face recognition (97), and reduced brightness perception was reported in one subject with developmental prosopagnosia (98). Impaired face perception in Alzheimer’s disease has been attributed to reduced contrast sensitivity at low spatial frequencies (7), but one study of seven patients with acquired prosopagnosia found more consistent deficits at higher spatial frequencies (99). Curvature perception has also been studied. The face has a myriad of curved surfaces, and impaired curvature perception was reported in a subject with developmental prosopagnosia (100, 101). However, in the series of subjects with acquired prosopagnosia, a deficit in curvature perception was found in only one subject, who also had the most difficulty on tests of basic-level object recognition (99).




Figure 4. Axial images from magnetic resonance imaging of two patients with ‘associative’ prosopagnosia. A) Unilateral anterior temporal damage in a 40 year-old man (R-AT3) presenting with prosopagnosia. B) Bilateral anterior temporal damage in a 47 year-old woman (B-AT2) presenting with prosopagnosia.

The problems with judging the spatial relationship between features demonstrated in apperceptive prosopagnosia (83) has raised the question of whether this is a deficit in perceiving fine spatial structure that extends to other visual stimuli. Some studies have shown that patients with acquired prosopagnosia have problems perceiving the spatial relationships between dots arranged in face and non-face-like configurations (99). This deficit was specifically linked to judgments about object shape, as their perception of the spatial relationships between different objects was not affected (102). Interestingly, their difficulty with the spatial arrangement of dot patterns continued to show influences of whole-object perception (103). Their ability to perceive local spatial relationships improved when more dots were available, suggesting that they gained by being able to reference dots to multiple points in the whole arrangement, when size or orientation changed between stimuli, which emphasized referencing to the whole object, or when the global arrangement had a more regular structure. Similarly, it has been reported that some patients with acquired prosopagnosia show normal influences of global whole-object structure with Navon letters (103, 104). Thus, while their impairment with judging the fine spatial metrics between parts of an object may be impaired not just for faces but also for other stimuli (but see case GG for an exception; (105)), any impairment in holistic processing may be specific for faces. This may also be an important distinction from congenital prosopagnosia, where differences in holistic processing for non-face objects like Navon letters have been demonstrated

(106, 107), as discussed in the article by Avidan and Behrmann in this issue.

3.2. Associative/amnestic prosopagnosia

In this variant, face perception is relatively intact, but there is a failure of this perceptual information to gain access to face memory stores (face recognition units) (37, 45, 108). This type of prosopagnosia is associated with anterior temporal lesions (39, 109, 110), and may be due to either a disconnection between facial percepts and the memory stores (43, 57), or a loss of facial memories. Similar to apperceptive prosopagnosia, *associative/amnestic* prosopagnosia can occur following either right unilateral damage, or bilateral damage (Figure 4).

To demonstrate intact or relatively preserved face perception, one can use the Benton Face Recognition Test or the Cambridge Face Perception Test. It should be noted that experimental tests of face discrimination can still show some perceptual deficits that are generally milder than those seen in patients with occipitotemporal damage. For example, even though patients with anterior temporal lesions do better at perceiving the spatial relations between facial features, some may fail to integrate these spatial relations with each other across the entire face: again, a type of whole-face processing deficit (110). Also, tests with morphed faces show that at least some patients with anterior temporal lesions have difficulty appreciating subtle variations in facial shape (Figure 2).

Acquired prosopagnosia: structure and perception

To probe the status of face memory, one can use probes of facial imagery, where the subject has to answer questions based on what they can recall about famous faces, without actually seeing them (43, 109). Failure to do this better than chance suggests a severe loss of facial memories, and has been noted in prosopagnosic patients with anterior temporal lesions and relatively preserved facial perceptual skills, while patients with posterior occipitotemporal lobe lesions show only mild impairments, despite their much more degraded face perception (60). Nevertheless, the fact that even occipitotemporal lesions do cause milder impairments in face imagery reinforces the point that the difference between these two groups of patients is often a relative rather than an absolute dichotomy. Hence prosopagnosic subjects with anterior temporal lesions are relatively better at facial perception and relatively worse on tests of facial imagery, compared to prosopagnosic patients with occipitotemporal lesions (38).

It has been argued by some that loss of facial memories should be considered an ‘*amnesic*’ deficit and distinct from inability to access facial memories through disconnection, with the term ‘*associative*’ reserved for the latter (15). In common, both amnesic and associative variants would show intact performance on encoding of facial structure on experimental tests or clinical assessments like the Benton Face Recognition Test or Cambridge Face Perception Test, and show no recognition or sense of familiarity on tests like the Warrington Recognition Memory Test or the Cambridge Face Memory Test. However, facial memories would still be intact in the disconnection scenario, and so these subjects should still perform well on tests of facial imagery, while those with the amnesic variant would fail (57). At this point, there are no known patients with impaired familiarity but both preservation of perceptual encoding of faces and also normal facial imagery: hence the pure associative non-amnesic variant remains a theoretical construct. For this reason, we tend to refer to prosopagnosic patients with preserved perceptual encoding and impaired facial imagery as having an *associative* variant, or sometimes an *associative/amnesic* variant, to stress the loss of facial memories.

While the classic disconnection proposed for associative prosopagnosia lies between perceptual processes and face memory stores in the face recognition units, another disconnection between face memories and person-identity nodes may also occur. This allows the patient to recognize that the face is familiar, because of a successful match between the percept and the correct facial memory, but they fail to retrieve the name or any biographical information about that person, because of failure to activate the corresponding person-identity node, even though name and biographical information can be accessed through other routes such as voice. Hence this condition may be better termed “*prosop-anomia*”, and has been described in a patient with an unusual left hemispheric lesion (111) and in another with bilateral temporal damage (47). Of note, in the first case, the defect was not specific to faces, as the patient also had anomia for places and other objects. It is of interest to speculate how a prosop-anomic

patient would do on tests of facial imagery: if the top-down connection between person-identity node and facial recognition units is also severed, then they should have as much difficulty in conjuring up the face to a name as do patients with the *associative/amnesic* variant of prosopagnosia.

As a last point, it is important to distinguish associative prosopagnosia from *people-specific amnesia*. Unlike associative prosopagnosic subjects who can still recognize people from other sensory cues, damage to the person identity nodes can result in the inability to recollect other people via any cues, including spoken names or voices. Although these patients present with a memory deficit specific to people with other types of memories intact, these patients are not prosopagnosic. People-specific amnesia has been described in patients with right temporal pole lesions (39, 40, 112, 113), and is consistent with neuroimaging studies showing that both name and face recognition activate the anterior middle temporal gyrus and temporal pole (114, 115). Although many prosopagnosic subjects claim to recognize others by voice, this has seldom been verified by formal tests, which may be particularly important in those with anterior temporal lesions.

3.3. Impairments of other functions

In addition to their difficulties with face recognition, prosopagnosic subjects often have other deficits, most often because naturally occurring damage is not limited to the face network but also involves neighboring structures. Damage to the optic radiations or striate cortex is common in those with occipitotemporal lesions, and causes visual field deficits, often in the left or bilateral upper quadrants, or sometimes a left hemianopia (2, 43, 99). Additional damage to the lingual and medial fusiform gyri causes achromatopsia. Also, many prosopagnosic patients with occipito-temporal damage have topographagnosia, or difficulty in navigating in familiar surroundings (20, 39, 41, 48, 53, 116, 117). Functional imaging studies show that buildings and places activate a specific region in occipito-temporal cortex, the ‘*parahippocampal place area*’, which is adjacent to the fusiform face area (118). Lesions here may cause a form of topographagnosia related to impaired recognition of landmarks. Field defects and problems with colour vision or navigation are less likely in prosopagnosic patients with anterior temporal damage, but these subjects may have minor visual or verbal memory disturbances (20, 119).

3.4. Face specificity versus expertise

Related to the issue of associated deficits is the controversy about whether prosopagnosia is a truly face-specific deficit caused by damage to a module or network dedicated to the processing of faces alone (120), or if this reflects damage to an expertise network, required for making subtle differentiations between similar exemplars of the same object category, of which faces are merely the most dramatic and universal example (121). Although some prosopagnosic subjects may also have a mild visual object agnosia (41, 48), it is generally agreed that they can identify objects at a basic level or category, for example being able to distinguish faces from cars, and from flowers.

Acquired prosopagnosia: structure and perception

What is argued is how well they do at more subtle distinctions, particularly within object categories, for example, distinguishing between types of cars or species of flowers. In support of face modularity are those reports of prosopagnosic subjects who can still identify personal belongings (122), individual animals (52, 123), specific places (39, 52), cars (52, 124, 125), flowers (39), vegetables (124, 126), and eyeglasses (127). In support of the expertise hypothesis are reports of prosopagnosic subjects who cannot identify types ('subordinate categories') of cars, food, or coins, or specific unique exemplars of buildings, handwriting, or personal clothing (17, 48, 128, 129).

There are many logical difficulties in settling this issue. On the one hand, impairments with recognizing other types of objects may simply be due to damage to other visual association cortex adjacent to face-processing regions, as is the case for achromatopsia and topographagnosia. On the other hand, claims of preserved recognition for another object class may be met with objections that either testing did not include a sufficient range of other object categories or else it did not test in sufficient detail. For example, some prosopagnosic subjects can show apparently normal accuracy rates on tests of their recognition of other objects, but analysis of reaction times or signal detection measures may reveal an impairment (130). Finally, a significant issue is that, while it is reasonable to assume nearly universal expertise for faces among human subjects, the same assumption cannot be made for most other visual objects. Hence, to evaluate the expertise hypothesis correctly, one must take into account how proficient a prosopagnosic subject was with the object type being tested before the onset of their problem (131). For example, the average person should be able to tell a hawk from an eagle, but a bird-watcher should be able to do better, perhaps distinguishing the *buteo* from *accipiter* genera of hawks. Showing intact hawk/eagle discrimination on a test may be evidence for preserved bird discrimination in the average person, but if this was the best discrimination a prosopagnosic bird-watcher could achieve after their lesion, one would suspect that they had impaired bird recognition as well.

In most situations, there is no opportunity to assess pre-morbid visual expertise with objects prior to the onset of acquired prosopagnosia. To deal with this difficulty, one might try to find a category for which subject interest and experience may be nearly as universal as with faces: this was the motivation for one study that found that most prosopagnosic subjects were also impaired in vegetable and fruit recognition (60, 132). However, for most object categories a degree of expertise homogenous across the population is unlikely and varying expertise will almost certainly be reflected in performance on tests. Even for food items, one might expect that a professional chef would be better than the average person. What is needed is some surrogate post-lesion measure that is highly likely to be correlated with pre-morbid visual expertise for other objects. Based on the assumption that semantic processing should be unaffected in a perceptual disorder such as prosopagnosia, one study examined the relationship

between verbal semantic knowledge and visual recognition of cars. These two measures are highly correlated in healthy subjects (133). This tight relationship allows one to predict from a prosopagnosic subject's semantic score what their visual recognition score should be if other object processing was indeed preserved. Based on this adjustment for pre-morbid expertise, our updated results in a group of 10 prosopagnosic subjects show that most subjects fell below the visual performance predicted by their verbal semantic score (Figure 5). In fact, convincing evidence for spared visual car recognition was found in only one subject, who had a right anterior temporal lesion from herpes encephalitis (Figure 4A).

3.5. Covert face recognition

Despite the professed inability of prosopagnosic subjects to recognize faces, in some there remains an unconscious or 'covert' face recognition (134-136). Covert face familiarity in prosopagnosia has been demonstrated with a wide variety of techniques, for which a taxonomy has been proposed (136). There are psychophysiologic measures such as electrodermal skin conductance (45, 117, 119), and electrophysiological measures such as visual evoked potentials (137). Behavioural methods can be direct, as with name-cued forced choice guessing of identity (12, 58, 138), learning of paired face-name associations (52, 58, 138, 139), and eye movements when viewing famous versus anonymous faces (92, 140). Finally, indirect behavioural techniques seek to determine if faces that subjects do not recognize overtly nevertheless influence the way subjects perform other tasks that do not involve recognizing the identity of the face. Classic examples include studies of priming and interference effects of faces on tasks involving names, such as sorting famous names according to occupation (141-143), or indicating whether the names are familiar or not (139). These studies show that whether the face matches the name in the task can influence the speed of performance, even though the subjects do not recognize the face.

The mechanisms responsible for covert recognition continue to be debated. One possibility, the dissociated dual-pathway hypothesis, is based on disconnection models, and proposed that covert recognition represents surviving processing in an intact dorsal occipitoparietal pathway to the amygdala that parallels the damaged occipitotemporal structures that support explicit recognition (144). Another possibility, the residual network activity hypothesis, is that it represents residual weak function of a damaged face-processing network, as supported by computer simulations (145-147), as well as observations that covert priming effects are correlated with the degree of residual overt recognition in prosopagnosia (143), a finding that is predicted by the residual network activity hypothesis but not by the dissociated dual-pathway hypothesis.

4. DEVELOPMENTAL PROSOPAGNOSIA

Before concluding, it is worth devoting a few words to developmental prosopagnosia, a condition that has received increasing attention since its first report in 1976




Figure 5. Car recognition in prosopagnosia, adjusted for semantic knowledge as an index of pre-morbid expertise. Visual recognition (naming the manufacturer of a car whose image is shown on the screen) is plotted as a function of verbal semantic knowledge (naming the manufacturer of a car whose name is given, without showing its image). In healthy controls the visual and verbal scores are highly correlated (solid line shows linear regression). The scores for 10 prosopagnosic patients are shown (blue = right anterior temporal, green = bilateral anterior temporal, red = right occipitotemporal, orange = bilateral occipitotemporal). The relationship between visual and verbal scores is lower for prosopagnosic subjects (dashed line) than it is for controls (solid line), indicating that their visual recognition of cars is worse than predicted by their verbal semantic knowledge. Only one prosopagnosic subject (RAT3) has convincing evidence of intact car recognition.

(148), in particular regarding its parallels with acquired prosopagnosia. Again, it is important first to draw some distinctions within the variety of face processing deficits that may originate in early life. In some cases prosopagnosia can be acquired because of brain injury during early childhood, when face expertise is still developing, while in other cases there is visual evidence of a cerebral malformation or an *in utero* insult (149). Also, impaired face recognition may be part of another congenital disorder, such as autism or Asperger syndrome (150). However, the term developmental prosopagnosia is best preserved for those patients in whom impaired face recognition is the chief or sole deficit, and in whom imaging does not reveal any evidence of a structural lesion: other cases may be referred to as early-onset acquired prosopagnosia.

Previous studies have estimated that up to 2-2.9% of the population may have developmental prosopagnosia (151, 152). However, when the diagnosis is based on the simple statistical criterion that a subject's score on a face recognition test lies more than 2 standard deviations below the mean (151), this estimate is tautological. (If the scores of a healthy population follow a normal distribution, 2.5% of its members will by definition score worse than 2 standard deviations below the mean.) Hence these figures are almost certainly an over-estimation of the true incidence of developmental prosopagnosia. Such liberal diagnostic criteria confound healthy subjects who are just worse than average with faces and those with a truly pathological problem.

Nevertheless it is undoubtedly true that there are some subjects born with extremely poor face recognition, which may reflect anomalous cerebral development. In some cases there may be a genetic contribution, as some individuals have family members with similar face recognition problems (107, 153, 154). This has led to the suggestion that developmental prosopagnosia may be an autosomal dominant trait (152). However, here again having several family members with poor face recognition is not necessarily evidence of a pathologic entity. Normal variation in face recognition abilities may also have a genetic component, as shown by twin studies that show greater correlation between face recognition abilities in monozygotic than in dizygotic twins (155). Besides genetic defects, other factors such as the failure to orient towards faces in early life (156), and certain personality traits such as shyness (157), may influence the development of face recognition abilities, given that these continue to evolve during early life.

The structural and functional parallels between acquired and developmental prosopagnosia are of interest. Neuroimaging studies have suggested a variety of subtler structural, functional, and connectivity anomalies in developmental prosopagnosia, with no consensus yet. Some studies have found reduced activity or face-selectivity in components of the core face network (106, 158-160), while others found relatively normal activation (161-165). Despite normal activation, one study did find reduced grey matter volume in portions of the inferior temporal cortex associated with face processing, which correlated with

facial identification performance (166). Similarly, the degree of face-selectivity of the right fusiform face area in developmental prosopagnosia has been shown to correlate with performance on face identification measures (160). Others suggest that the abnormality in developmental prosopagnosia lies not in the core face network but beyond it. A reduction in grey matter volume in the right anterior fusiform gyrus (anterior to the fusiform face area) and a corresponding increase in the right middle temporal gyrus, has been reported in developmental prosopagnosia, with this reduction correlating with face recognition deficits (167). Reduced structural connectivity in occipitotemporal cortex has also been shown in developmental prosopagnosia (168), and has been followed by demonstrations of reduced functional connectivity between the core face network and anterior temporal regions involved in processing identity (165).

Of course, it is not necessarily the case that all subjects with developmental prosopagnosia have the same structural anomaly, given the variety of lesions in the acquired form. It may be that some subjects have reduced activity of the core face-network, while others have reduced connectivity from this network to anterior temporal structures, perhaps due to different genetic defects. If so, this may also be paralleled by functional subtypes just as found in acquired prosopagnosia. However, at present the question of functional subtypes of developmental prosopagnosia has only recently begun to be addressed (169, 170), and further confirmatory studies of larger series of subjects with uniform test batteries of diverse face processing functions would be welcome. See Avidan and Behrmann (this issue) for more discussion on this topic.

5. SUMMARY

Prosopagnosia is a family of disorders that differ in the responsible cognitive deficit and accompanying structural lesions. In the acquired form, difficulties perceiving facial structure is characteristic of apperceptive prosopagnosia, which is associated with right or bilateral occipitotemporal lesions that involve the fusiform and/or occipital face areas. The perceptual problem may lie in extracting the fine spatial metrics of the complex shape that is the face, which is usually done rapidly in a holistic fashion. Whether this type of perceptual analysis is specific to faces or affects other object categories continues to be debated: current evidence suggests that some impairment for other object classes, particularly those for which the subject has some expertise, may be the rule rather than the exception. Difficulties accessing facial memories because of disconnection or loss of those memories is typical of associative/amnesic prosopagnosia, which involves more anterior structures in the right or both temporal lobes. As mentioned, most of the preceding points have been learned from cases of acquired prosopagnosia. Developmental prosopagnosia is a specific deficit that is present from birth, with evidence for some genetic contribution. Whether there are similar structural and functional parallels between acquired and developmental prosopagnosia is not yet known.

6. ACKNOWLEDGMENTS

This work was supported by CIHR grants MOP-102567 and MOP-106511. JB was supported by a Canada Research Chair.

7. REFERENCES

1. J. Bodamer: Prosopagnosie. *Arch Psychiatr Nervenkr*, 179, 6-54 (1947)
2. J. Barton: Disorders of face perception and recognition. *Neurologic Clinics*, 21, 521-548 (2003)
3. S. Rapcsak, M. Polster, M. Glisky and J. Comer: False recognition of unfamiliar faces following right hemisphere damage: neuropsychological and anatomical observations. *Cortex*, 32, 593-611 (1996)
4. L. Tejeria, R. A. Harper, P. H. Artes and C. M. Dickinson: Face recognition in age related macular degeneration: perceived disability, measured disability, and performance with a bioptic device. *Br J Ophthalmol*, 86(9), 1019-26 (2002)
5. M. Mendez, R. Martin, K. Smyth and P. Whitehouse: Disturbances of person identification in Alzheimer's disease. A retrospective study. *J Nerv Ment Dis*, 180, 94-6 (1992)
6. M. Roudier, P. Marcie, A. Grancher, C. Tzortzis, S. Starkstein and F. Boller: Discrimination of facial identity and of emotions in Alzheimer's disease. *J Neurol Sci*, 154, 151-8 (1998)
7. A. Cronin-Coulomb, M. Cronin-Coulomb, T. Dunne, A. Brown, K. Jain, P. Cipolloni and S. Auerbach: Facial frequency manipulation normalizes face discrimination in AD. *Neurology*, 54, 2316-8 (2000)
8. T. S. Lim, H. Y. Lee, J. J. Barton and S. Y. Moon: Deficits in face perception in the amnesic form of mild cognitive impairment. *J Neurol Sci*, 309(1-2), 123-7 (2011)
9. A. Janati: Kluver-Bucy syndrome in Huntington's chorea. *J Nerv Ment Dis*, 173, 632-5 (1985)
10. H. Dewick, J. Hanley, A. Davies, J. Playfer and C. Turnbull: Perception and memory for faces in Parkinson's disease. *Neuropsychologia*, 29, 785-802 (1991)
11. R. Cousins, J. R. Hanley, A. D. Davies, C. J. Turnbull and J. R. Playfer: Understanding memory for faces in Parkinson's disease: the role of configural processing. *Neuropsychologia*, 38(6), 837-47 (2000)
12. J. J. Barton, M. Cherkasova and M. O'Connor: Covert recognition in acquired and developmental prosopagnosia. *Neurology*, 57(7), 1161-8 (2001)
13. E. Warrington: Warrington Recognition Memory Test. Western Psychological Services, Los Angeles (1984)

Acquired prosopagnosia: structure and perception

14. B. Duchaine and K. Nakayama: Dissociations of face and object recognition in developmental prosopagnosia. *J Cogn Neurosci*, 17, 249-61 (2005)
15. A. Damasio, D. Tranel and H. Damasio: Face agnosia and the neural substrates of memory. *Ann Rev Neurosci*, 13, 89-109 (1990)
16. J. Meadows: The anatomical basis of prosopagnosia. *J Neurol Neurosurg Psychiatr*, 37, 489-501 (1974)
17. A. Damasio, H. Damasio and G. van Hoessen: Prosopagnosia: anatomic basis and behavioral mechanisms. *Neurology*, 32, 331-41 (1982)
18. H. Hécaen and R. Angelergues: Agnosia for faces (prosopagnosia). *Arch Neurol*, 7, 92-100 (1962)
19. E. de Renzi: Prosopagnosia in two patients with CT scan evidence of damage confined to the right hemisphere. *Neuropsychologia*, 24(3), 385-389 (1986)
20. T. Landis, J. Cummings, L. Christen, J. Bogen and H.-G. Imhof: Are unilateral right posterior lesions sufficient to cause prosopagnosia ? Clinical and radiological findings in six additional patients. *Cortex*, 22, 243-52 (1986)
21. A. Tzavaras, L. Merienne and M. C. Masure: Prosopagnosia, amnesia and language disorders caused by left temporal lobe injury in a left-handed man. *L'Encéphale*, 62(4), 382-394 (1973)
22. M. Eimer and R. A. McCarthy: Prosopagnosia and structural encoding of faces : Evidence from event-related potentials. *Neuroreport*, 10(2), 255-259 (1999)
23. A. J. Mattson, H. S. Levin and J. Grafman: A Case of Prosopagnosia Following Moderate Closed Head Injury with Left Hemisphere Focal Lesion. *Cortex*, 36(1), 125-137 (2000)
24. J. J. Barton: Prosopagnosia associated with a left occipitotemporal lesion. *Neuropsychologia* (2008)
25. N. Kanwisher, J. McDermott and M. Chun: The fusiform face area: a module in human extrastriate cortex specialized for face perception. *J Neurosci*, 17, 4302-11 (1997)
26. J. J. Barton, C. J. Fox, A. Sekunova and G. Iaria: Encoding in the Visual Word Form Area: An fMRI Adaptation Study of Words versus Handwriting. *J Cogn Neurosci* (2009)
27. B. Rossion, L. Dricot, A. Devolder, J.-M. Bodart, M. Crommelinck, B. d. Gelder and R. Zoontjes: Hemispheric Asymmetries for Whole-Based and Part-Based Face Processing in the Human Fusiform Gyrus. *Journal of Cognitive Neuroscience*, 12(5), 793-802 (2000)
28. T. Ettlin, M. Beckson, D. Benson, J. Langfitt, E. Amos and G. Pineda: Prosopagnosia: a bihemispheric disorder. *Cortex*, 28, 129-34 (1992)
29. V. Bruce and A. Young: Understanding face recognition. *British Journal of Psychology*, 77(3), 305-327 (1986)
30. D. Perrett, J. Hietanen, M. Oram and P. Benson: Organization and functions of cells responsive to faces in the temporal cortex. *Phil Trans R Soc Lond B*, 335, 23-30 (1992)
31. I. Gauthier and N. Logothetis: Is face recognition not so unique after all? *Cognitive Neuropsychol*, 17, 125-142 (2000)
32. J. V. Haxby, M. I. Gobbini, M. L. Furey, A. Ishai, J. L. Schouten and P. Pietrini: Distributed and overlapping representations of faces and objects in ventral temporal cortex. *Science*, 293(5539), 2425-30 (2001)
33. A. J. Calder and A. W. Young: Understanding the recognition of facial identity and facial expression. *Nat Rev Neurosci*, 6(8), 641-651 (2005)
34. C. J. Fox, H. M. Hanif, G. Iaria, B. C. Duchaine and J. J. Barton: Perceptual and anatomic patterns of selective deficits in facial identity and expression processing. *Neuropsychologia*, 49(12), 3188-200 (2011)
35. V. Bruce and A. Young: Understanding face recognition. *Brit J Psychol*, 77, 305-27 (1986)
36. A. Burton, V. Bruce and P. Hancock: From pixels to people: a model of familiar face recognition. *Cognitive Sci*, 23, 1-31 (1999)
37. E. de Renzi, P. Faglioni, D. Grossi and P. Nichelli: Apperceptive and associative forms of prosopagnosia. *Cortex*, 27, 213-21 (1991)
38. J. J. S. Barton: Structure and function in acquired prosopagnosia: Lessons from a series of 10 patients with brain damage. *Journal of Neuropsychology*, 2(1), 197-225 (2008)
39. J. Evans, A. Heggs, N. Antoun and J. Hodges: Progressive prosopagnosia associated with selective right temporal lobe atrophy. *Brain*, 118, 1-13 (1995)
40. G. Gainotti, M. Ferraccioli, D. Quaranta and C. Marra: Cross-modal recognition disorders for persons and other unique entities in a patient with right fronto-temporal degeneration. *Cortex*, 44(3), 238-48 (2008)
41. A. Young and H. Ellis: Childhood prosopagnosia. *Brain and Cognition*, 9, 16-47 (1989)
42. I. Kracke: Developmental prosopagnosia in Asperger syndrome: presentation and discussion of an individual case. *Dev Med Child Neurol*, 36, 873-86 (1994)
43. N. Takahashi, M. Kawamura, K. Hirayama, J. Shiota and O. Isono: Prosopagnosia: a clinical and anatomic study of four patients. *Cortex*, 31, 317-29 (1995)

Acquired prosopagnosia: structure and perception

44. J. J. Barton: Disorder of higher visual function. *Curr Opin Neurol*, 24(1), 1-5 (2011)
45. D. Tranel and A. Damasio: Knowledge without awareness: an autonomic index of facial recognition by prosopagnosics. *Science*, 228, 1453-1454 (1985)
46. A. Young, J. Aggleton, D. Hellawell, M. Johnson, P. Broks and J. Hanley: Face processing impairments after amygdalotomy. *Brain*, 118, 15-24 (1995)
47. L. J. Tippet, L. A. Miller and M. J. Farah: Prosopamnesia: a selective impairment in face learning. *Cogn Neuropsychol*, 17(1), 241-55 (2000)
48. E. de Haan and R. Campbell: A fifteen year follow-up of a case of developmental prosopagnosia. *Cortex*, 27, 489-509 (1991)
49. R. Campbell, C. Heywood, A. Cowey, M. Regard and T. Landis: Sensitivity to eye gaze in prosopagnosic patients and monkeys with superior temporal sulcus ablation. *Neuropsychologia*, 28, 1123-42 (1990)
50. K. Humphreys, G. Avidan and M. Behrmann: A detailed investigation of facial expression processing in congenital prosopagnosia as compared to acquired prosopagnosia. *Exp Brain Res*, 176(2), 356-73 (2007)
51. B. C. Stephan, N. Breen and D. Caine: The recognition of emotional expression in prosopagnosia: decoding whole and part faces. *J Int Neuropsychol Soc*, 12(6), 884-95 (2006)
52. R. Bruyer, C. Laterre, X. Seron, P. Feyereisen, E. Strypstein, E. Pierrard and D. Rectem: A case of prosopagnosia with some preserved covert remembrance of familiar faces. *Brain and Cognition*, 2, 257-84 (1983)
53. J. Sergent and J.-G. Villemure: Prosopagnosia in a right hemispherectomized patient. *Brain*, 112, 975-95 (1989)
54. J. Sergent and M. Poncet: From covert to overt recognition of faces in a prosopagnosic patient. *Brain*, 113, 989-1004 (1990)
55. D. Tranel, A. R. Damasio and H. Damasio: Intact recognition of facial expression, gender, and age in patients with impaired recognition of face identity. *Neurology*, 38(5), 690-6 (1988)
56. C. J. Fox, G. Iaria and J. J. S. Barton: Defining the face processing network: Optimization of the functional localizer in fMRI. *Human Brain Mapping*, 30(5), 1637-1651 (2009)
57. C. J. Fox, G. Iaria and J. J. Barton: Disconnection in prosopagnosia and face processing. *Cortex*, 44(8), 996-1009 (2008)
58. J. Sergent and J.-L. Signoret: Implicit access to knowledge derived from unrecognized faces. *Cerebral Cortex*, 2, 389-400 (1992)
59. J. J. Barton, D. Z. Press, J. P. Keenan and M. O'Connor: Lesions of the fusiform face area impair perception of facial configuration in prosopagnosia. *Neurology*, 58(1), 71-8 (2002)
60. J. J. S. Barton: Structure and function in acquired prosopagnosia: lessons from a series of ten patients with brain damage. *J Neuropsychology*, 2, 197-225 (2008)
61. S. E. Bouvier and S. A. Engel: Behavioral deficits and cortical damage loci in cerebral achromatopsia. *Cereb Cortex*, 16, 183-91 (2006)
62. B. Rossion, R. Caldara, M. Seghier, A. M. Schuller, F. Lazeyras and E. Mayer: A network of occipito-temporal face-sensitive areas besides the right middle fusiform gyrus is necessary for normal face processing. *Brain*, 126(Pt 11), 2381-95 (2003)
63. C. Schiltz, B. Sorger, R. Caldara, F. Ahmed, E. Mayer, R. Goebel and B. Rossion: Impaired face discrimination in acquired prosopagnosia is associated with abnormal response to individual faces in the right middle fusiform gyrus. *Cereb Cortex*, 16(4), 574-86 (2006)
64. A. Benton and M. van Allen: Prosopagnosia and facial discrimination. *J Neurol Sci*, 15, 167-72 (1972)
65. E. de Renzi, P. Faglioni and H. Spinnler: The performance of patients with unilateral brain damage on face recognition tasks. *Cortex*, 4, 17-43 (1968)
66. G. Carlesimo and C. Caltagirone: Components in the visual processing of known and unknown faces. *J Clin Exp Neuropsychol*, 17, 691-705 (1995)
67. F. Parry, A. Young, J. Saul and A. Moss: Dissociable face processing impairments after brain injury. *J Clin Exp Neuropsychol*, 13, 545-58 (1991)
68. M. J. Farah: Visual Agnosia: disorders of visual recognition and what they tell us about normal vision. MIT Press, Cambridge (1990)
69. B. Duchaine, L. Germine and K. Nakayama: Family resemblance: Ten family members with prosopagnosia and within-class object agnosia. *Cognitive Neuropsychology*, 24(4), 419-430 (2007)
70. G. Malcolm, C. Leung and J. Barton: Regional variation in the inversion effect for faces: different patterns for feature shape, spatial relations, and external contour. *Perception*, 33, 1221-31 (2005)
71. K. Tanaka: Neuronal mechanisms of object recognition. *Science*, 262(5134), 685-688 (1993)
72. J. Tanaka and J. Sengco: Features and their configuration in face recognition. *Memory and Cognition*, 25, 583-92 (1997)
73. M. Farah, K. Wilson, H. Drain and J. Tanaka: The inverted face inversion effect in prosopagnosia: evidence

Acquired prosopagnosia: structure and perception

for mandatory, face-specific perceptual mechanisms. *Vision Res*, 35, 2089-93 (1995)

74. A. Young, D. Hellawell and D. Hay: Configurational information in face perception. *Perception*, 16, 747-59 (1987)

75. J. Sergent and J.-G. Villemure: Prosopagnosia in a Right Hemispherectomized Patient. *Brain*, 112(4), 975-995 (1989)

76. M. Ramon, T. Busigny and B. Rossion: Impaired holistic processing of unfamiliar individual faces in acquired prosopagnosia. *Neuropsychologia*, 48(4), 933-44 (2010)

77. G. Van Belle, P. De Graef, K. Verfaillie, T. Busigny and B. Rossion: Whole not hole: expert face recognition requires holistic perception. *Neuropsychologia*, 48(9), 2620-9 (2010)

78. G. Rhodes: Looking at faces: first-order and second-order features as determinants of facial appearance. *Perception*, 17, 43-63 (1988)

79. H. Leder and V. Bruce: When inverted faces are recognized: the role of configural information in face recognition. *Q J Exp Psychol A*, 53(2), 513-36 (2000)

80. J. Barton, J. Keenan and T. Bass: Discrimination of spatial relations and features in faces: effects of inversion and viewing duration. *Brit J Psychol*, 92, 527-49 (2001)

81. S. Yamane, S. Kaji and K. Kawano: What facial features activate face neurons in the inferotemporal cortex of the monkey? *Experimental Brain Research*, 73(1), 209-214 (1988)

82. M. Young and S. Yamane: Sparse population coding of faces in the inferotemporal cortex. *Science*, 256(5061), 1327-1331 (1992)

83. J. Barton, D. Press, J. Keenan and M. O'Connor: Lesions of the fusiform face area impair perception of facial configuration in prosopagnosia. *Neurology*, 58, 71-8 (2002)

84. S. Joubert, O. Felician, E. Barbeau, A. Sontheimer, J. Barton, M. Ceccaldi and M. Poncet: Impaired configurational processing in a case of progressive prosopagnosia associated with right temporal lobe atrophy. *Brain*, 126, 2537-2550 (2003)

85. J. Shepherd, G. Davies and H. Ellis: Studies of cue saliency. In: *Perceiving and remembering faces*. Ed G. Davies, H. Ellis & J. Shepherd. Academic Press, London (1981)

86. G. L. Malcolm, L. J. Lanyon, A. J. Fugard and J. J. Barton: Scan patterns during the processing of facial expression versus identity: an exploration of task-driven and stimulus-driven effects. *J Vis*, 8(8), 2 1-9 (2008)

87. P. Schyns, L. Bonnar and F. Gosselin: Show me the features! Understanding recognition from the use of visual information. *Psychol Sci*, 13, 402-9 (2002)

88. C. Vinette, Gosselin, F., Schyns, P.G.: Spatio-temporal dynamics of face recognition in a flash: it's in the eyes. *Cognitive Science*, 28(2), 289-301 (2004)

89. M. L. Smith, G. W. Cottrell, F. Gosselin and P. G. Schyns: Transmitting and decoding facial expressions. *Psychol Sci*, 16(3), 184-189 (2005)

90. N. Haig: How faces differ - a new comparative technique. *Perception*, 14, 601-15 (1985)

91. N. Haig: Investigating face recognition with an image processing computer. In: *Aspects of Face Processing*. Ed J. M. Ellis HD, Newcombe F, Young A. Martinus Nejhoff, Dordrecht (1986)

92. J. Barton, N. Radcliffe, M. Cherkasova and J. Edelman: Scan patterns during the processing of facial identity in prosopagnosia. *Experimental Brain Research*, 181(2), 199-211 (2007)

93. J. J. S. Barton, D. Z. Press, J. P. Keenan and M. O'Connor: Lesions of the fusiform face area impair perception of facial configuration in prosopagnosia. *Neurology*, 58(1), 71-78 (2002)

94. R. Caldara, P. Schyns, E. Mayer, M. Smith, F. Gosselin and B. Rossion: Does prosopagnosia take the eyes out of face representations? Evidence for a defect in representing diagnostic facial information following brain damage. *J Cogn Neurosci*. (2005)

95. C. M. Bukach, D. N. Bub, I. Gauthier and M. J. Tarr: Perceptual expertise effects are not all or none: spatially limited perceptual expertise for faces in a case of prosopagnosia. *J Cogn Neurosci*, 18(1), 48-63 (2006)

96. C. M. Bukach, R. Le Grand, M. D. Kaiser, D. N. Bub and J. W. Tanaka: Preservation of mouth region processing in two cases of prosopagnosia. *J Neuropsychol*, 2(Pt 1), 227-44 (2008)

97. R. Kemp, G. Pike, P. White and A. Musselman: Perception and recognition of normal and negative faces - the role of shape from shading and pigmentation cues. *Perception*, 25, 37-52 (1996)

98. R. Campbell: Face to face: interpreting a case of developmental prosopagnosia. In: *Mental Lives. Case studies in cognition*. Ed R. Campbell. Blackwell, Oxford, UK (1992)

99. J. Barton, M. Cherkasova, D. Press, J. Intriligator and M. O'Connor: Perceptual function in prosopagnosia. *Perception*, 33, 939-956 (2004)

100. S. Kosslyn, S. Hamilton and J. Bernstein: The perception of curvature can be selectively disrupted in prosopagnosia. *Brain Cognition*, 27, 36-58 (1995)

Acquired prosopagnosia: structure and perception

101. B. Laeng and V. S. Caviness: Prosopagnosia as a deficit in encoding curved surface. *J Cogn Neurosci*, 13(5), 556-76 (2001)
102. J. J. Barton, G. L. Malcolm and R. L. Hefter: Spatial processing in Balint syndrome and prosopagnosia: a study of three patients. *J Neuroophthalmol*, 27(4), 268-74 (2007)
103. J. J. S. Barton: What is meant by impaired configural processing in acquired prosopagnosia? . *Perception* 39, 242-60 (2009)
104. T. Busigny and B. Rossion: Holistic processing impairment can be restricted to faces in acquired prosopagnosia: Evidence from the global/local Navon effect. *J Neuropsychol* (2010)
105. T. Busigny, S. Joubert, O. Felician, M. Ceccaldi and B. Rossion: Holistic perception of the individual face is specific and necessary: Evidence from an extensive case study of acquired prosopagnosia. *Neuropsychologia*, 48(14), 4057-4092 (2010)
106. S. Bentin, J. M. DeGutis, M. d'Esposito and L. C. Robertson: Too many trees to see the forest: performance, event-related potential, and functional magnetic resonance imaging manifestations of integrative congenital prosopagnosia. *J Cogn Neurosci*, 19(1), 132-146 (2007)
107. M. Behrmann, G. Avidan, J. Marotta and R. Kimchi: Detailed exploration of face-related processing in congenital prosopagnosia: 1. Behavioral findings. *J Cogn Neurosci*, 17, 1130-49 (2005)
108. A. Damasio, D. Tranel and H. Damasio: Face agnosia and the neural substrates of memory. *Ann Rev Neurosci*, 13, 89-109 (1990)
109. J. J. Barton and M. Cherkasova: Face imagery and its relation to perception and covert recognition in prosopagnosia. *Neurology*, 61(2), 220-5 (2003)
110. J. J. Barton, J. Zhao and J. P. Keenan: Perception of global facial geometry in the inversion effect and prosopagnosia. *Neuropsychologia*, 41(12), 1703-11 (2003)
111. P. Verstichel and L. Chia: Difficulties in face identification after lesion in the left hemisphere. *Rev Neurol (Paris)*, 155(11), 937-43 (1999)
112. H. Ellis, A. Young and E. Critchley: Loss of memory for people following temporal lobe damage. *Brain*, 1989, 1469-83 (1989)
113. J. Hanley, A. Young and N. Pearson: Defective recognition of familiar people. *Cogn Neuropsychol*, 6, 179-210 (1989)
114. M. L. Gorno-Tempini and C. J. Price: Identification of famous faces and buildings: a functional neuroimaging study of semantically unique items. *Brain*, 124(Pt 10), 2087-97 (2001)
115. C. Leveroni, M. Seidenberg, A. Mayer, L. Mead, J. Binder and S. Rao: Neural systems underlying the recognition of familiar and newly learned faces. *J Neurosci*, 20, 878-86 (2000)
116. D. Malone, H. Morris, M. Kay and H. Levin: Prosopagnosia: a double dissociation between the recognition of familiar and unfamiliar faces. *J Neurol Neurosurg Psychiatr*, 45, 820-2 (1982)
117. R. Bauer: Autonomic recognition of names and faces in prosopagnosia: a neuropsychological application of the guilty knowledge test. *Neuropsychologia*, 22, 457-69 (1984)
118. K. M. O'Craven and N. Kanwisher: Mental imagery of faces and places activates corresponding stimulus- specific brain regions. *J Cogn Neurosci*, 12(6), 1013-23 (2000)
119. R. Bauer and M. Verfaellie: Electrodermal discrimination of familiar but not unfamiliar faces in prosopagnosia. *Brain and Cognition*, 8, 240-52 (1988)
120. N. Kanwisher: Domain specificity in face perception. *Nat Neurosci*, 3(8), 759-63 (2000)
121. M. J. Tarr and I. Gauthier: FFA: a flexible fusiform area for subordinate-level visual processing automatized by expertise. *Nat Neurosci*, 3(8), 764-9 (2000)
122. E. de Renzi: Prosopagnosia in two patients with CT scan evidence of damage confined to the right hemisphere. *Neuropsychologia*, 24, 385-9 (1986)
123. J. McNeil and E. Warrington: Prosopagnosia: a face-specific disorder. *Quart J Exp Psychol*, 46A, 1-10 (1993)
124. K. Henke, S. Schweinberger, A. Grigo, T. Klos and W. Sommer: Specificity of face recognition: recognition of exemplars of non-face objects in prosopagnosia. *Cortex*, 34, 289-96 (1998)
125. T. Busigny, M. Graf, E. Mayer and B. Rossion: Acquired prosopagnosia as a face-specific disorder: ruling out the general visual similarity account. *Neuropsychologia*, 48(7), 2051-67 (2010)
126. M. J. Riddoch, R. A. Johnston, R. M. Bracewell, L. Boutsen and G. W. Humphreys: Are faces special? A case of pure prosopagnosia. *Cogn Neuropsychol*, 25(1), 3-26 (2008)
127. M. Farah, K. Levinson and K. Klein: Face perception and within-category discrimination in prosopagnosia. *Neuropsychologia*, 33, 661-74 (1995)
128. F. Lhermitte, F. Chain, R. Escourolle, B. Ducarne and B. Pillon: Étude anatomo-clinique d'un cas de prosopagnosie. *Rev Neurol*, 126, 329-346 (1972)
129. A. Whiteley and E. Warrington: Prosopagnosia: a clinical, psychological, and anatomical study of three

Acquired prosopagnosia: structure and perception

- patients. *J Neurol Neurosurg Psychiatry*, 40, 395-403 (1977)
130. I. Gauthier, M. Behrmann and M. Tarr: Can face recognition really be dissociated from object recognition? *J Cogn Neurosci*, 11, 349-70 (1999)
131. R. Diamond and S. Carey: Why faces are and are not special: an effect of expertise. *J Exp Psychol: General*, 115, 107-17 (1986)
132. J. Barton, M. Cherkasova, D. Press, J. Intriligator and M. O'Connor: Perceptual function in prosopagnosia. *J Neuro-ophthalmology*, in press (2004)
133. J. J. Barton, H. Hanif and S. Ashraf: Relating visual to verbal semantic knowledge: the evaluation of object recognition in prosopagnosia. *Brain : a journal of neurology*, 132(Pt 12), 3456-66 (2009)
134. R. Bruyer: Covert facial recognition in prosopagnosia: a review. *Brain and Cognition*, 15, 223-35 (1991)
135. A. Young: Covert recognition. In: *The neuropsychology of high-level vision*. Ed M. Farah&G. Ratcliff. Lawrence Erlbaum Associates, Hillsdale (1994)
136. J. J. S. Barton: La reconnaissance implicite dans la prosopagnosie. In: *Traitement et reconnaissance des visages : du percept à la personne*. Ed E. Barbeau, O. Felician&S. Joubert. Solal, Marseille (2009)
137. B. Renault, J.-L. Signoret, B. DeBruille, F. Breton and F. Bolgert: Brain potentials reveal covert facial recognition in prosopagnosia. *Neuropsychologia*, 27, 905-12 (1989)
138. J. McNeil and E. Warrington: Prosopagnosia: a re-classification. *Quart J Exp Psychol*, 43A, 267-87 (1991)
139. S. Schweinberger, T. Klos and W. Sommer: Covert face recognition in prosopagnosia: a dissociable function? *Cortex*, 31, 517-29 (1995)
140. M. Rizzo, R. Hurtig and A. Damasio: The role of scanpaths in facial recognition and learning. *Ann Neurol*, 22, 41-5 (1987)
141. E. de Haan, A. Young and F. Newcombe: Faces interfere with name classification in a prosopagnosic patient. *Cortex*, 23, 309-16 (1987)
142. A. Young, D. Hellawell and E. de Haan: Cross-domain semantic priming in normal subjects and a prosopagnosic patient. *Quart J Exp Psychol*, 40A, 561-80 (1988)
143. J. Barton, M. Cherkasova and R. Hefter: The covert priming effect of faces in prosopagnosia. *Neurology*, 63, 2062-2068 (2004)
144. D. Tranel, H. Damasio and A. Damasio: Double dissociation between overt and covert face recognition. *J Cognitive Neurosci*, 7, 425-432 (1995)
145. M. Farah, R. O'Reilly and S. Vecera: Dissociated overt and covert recognition as an emergent property of a lesioned neural network. *Psychological Review*, 100, 571-88 (1993)
146. A. Young and A. Burton: Simulating face recognition: implications for modelling cognition. *Cognitive Neuropsychology*, 16, 1-48 (1999)
147. R. O'Reilly and M. Farah: Simulation and explanation in neuropsychology and beyond. *Cognitive Neuropsychology*, 16, 49-72 (1999)
148. H. McConachie: Developmental prosopagnosia, a single case report. *Cortex*, 12, 76-82 (1976)
149. J. Barton, M. Cherkasova, D. Press, J. Intriligator and M. O'Connor: Developmental prosopagnosia: a study of three patients. *Brain and Cognition*, 51, 12-30 (2003)
150. J. Barton, M. Cherkasova, R. Hefter, T. Cox, M. O'Connor and D. Manoach: Are patients with social developmental disorders prosopagnosic? Perceptual heterogeneity in the Asperger and socio-emotional processing disorders. *Brain*, 127, 1706-1716 (2004)
151. D. C. Bowles, E. McKone, A. Dawel, B. Duchaine, R. Palermo, L. Schmalzl, D. Rivolta, C. E. Wilson and G. Yovel: Diagnosing prosopagnosia: effects of ageing, sex, and participant-stimulus ethnic match on the Cambridge Face Memory Test and Cambridge Face Perception Test. *Cogn Neuropsychol*, 26(5), 423-55 (2009)
152. I. Kennerknecht, N. Plumpe, S. Edwards and R. Raman: Hereditary prosopagnosia (HPA): the first report outside the Caucasian population. *J Hum Genet*, 52(3), 230-6 (2007)
153. B. C. Duchaine, T. Nieminen-von Wendt, J. New and T. Kulomaki: Dissociations of visual recognition in a developmental agnosic: evidence for separate developmental processes. *Neurocase*, 9(5), 380-9 (2003)
154. A. L. Diaz: Do I know you? A case study of prosopagnosia (face blindness). *J Sch Nurs*, 24(5), 284-9 (2008)
155. J. B. Wilmer, L. Germine, C. F. Chabris, G. Chatterjee, M. Williams, E. Loken, K. Nakayama and B. Duchaine: Human face recognition ability is specific and highly heritable. *Proc Natl Acad Sci U S A*, 107(11), 5238-41 (2010)
156. J. Morton and M. H. Johnson: CONSPEC and CONLERN: a two-process theory of infant face recognition. *Psychol Rev*, 98(2), 164-81 (1991)
157. P. M. Brunet, C. J. Mondloch and L. A. Schmidt: Shy children are less sensitive to some cues to facial recognition. *Child Psychiatry Hum Dev*, 41(1), 1-14 (2010)
158. N. Hadjikhani and B. de Gelder: The neural basis of prosopagnosia - an fMRI study. *Human Brain Mapping*, 16, 176-182 (2002)

Acquired prosopagnosia: structure and perception

159. D. A. Minnebusch, B. Suchan, O. Köster and I. Daum: A bilateral occipitotemporal network mediates face perception. *Behavioural Brain Research*, 198(1), 179-185 (2009)

160. N. Furl, L. Garrido, R. J. Dolan, J. Driver and B. Duchaine: Fusiform gyrus face selectivity relates to individual differences in facial recognition ability. *J Cogn Neurosci*, 23(7), 1723-40 (2011)

161. U. Hasson, G. Avidan, L. Y. Deouell, S. Bentin and R. Malach: Face-selective activation in a congenital prosopagnosic subject. *J Cogn Neurosci*, 15(3), 419-31 (2003)

162. G. Avidan, U. Hasson, R. Malach and M. Behrmann: Detailed exploration of face-related processing in congenital prosopagnosia: 2. Functional neuroimaging findings. *J Cogn Neurosci*, 17, 1150-67 (2005)

163. J. M. DeGutis, S. Bentin, L. C. Robertson and M. D'Esposito: Functional plasticity in ventral temporal cortex following cognitive rehabilitation of a congenital prosopagnosic. *J Cogn Neurosci*, 19(11), 1790-802 (2007)

164. G. Avidan and M. Behrmann: Functional MRI reveals compromised neural integrity of the face processing network in congenital prosopagnosia. *Curr Biol*, 19(13), 1146-50 (2009)

165. G. Avidan, M. Tanzer, F. Hadj-Bouziane, N. Liu, L. G. Ungerleider and M. Behrmann: Selective Dissociation Between Core and Extended Regions of the Face Processing Network in Congenital Prosopagnosia. *Cereb Cortex* (2013)

166. L. Garrido, N. Furl, B. Draganski, N. Weiskopf, J. Stevens, G. C. Tan, J. Driver, R. J. Dolan and B. Duchaine: Voxel-based morphometry reveals reduced grey matter volume in the temporal cortex of developmental prosopagnosics. *Brain*, 132(Pt 12), 3443-55 (2009)

167. M. Behrmann, G. Avidan, F. Gao and S. Black: Structural Imaging Reveals Anatomical Alterations in Inferotemporal Cortex in Congenital Prosopagnosia. *Cereb Cortex*, epub (2007)

168. C. Thomas, G. Avidan, K. Humphreys, K. J. Jung, F. Gao and M. Behrmann: Reduced structural connectivity in ventral visual cortex in congenital prosopagnosia. *Nat Neurosci*, 12(1), 29-31 (2009)

169. K. A. Dalrymple and B. Duchaine: Varied behavioral profiles in children with developmental prosopagnosia reveal dissociations in the developing face processing system. In: *Vision Sciences Society. Journal of Vision*, Naples, Florida (2013)

170. R. Stollhoff, J. Jost, T. Elze and I. Kennerknecht: Deficits in long-term recognition memory reveal dissociated subtypes in congenital prosopagnosia. *PLoS One*, 6(1), e15702 (2011)

Key Words: Face Recognition, Perception, Face Imagery, Fusiform Gyrus, Anterior Temporal, Review

Send correspondence to: Jason J S Barton, Neuro-ophthalmology Section D, VGH Eye Care Center, 2550 Willow Street, Vancouver BC Canada V5Z 3N9, Tel: 604-875-4339, Tel: 604-875-4302, E-mail: jasonbarton@shaw.ca