

Neuroprotective agents and modulation of temporal lobe epilepsy

Maria Jose da Silva Fernandes¹, Jose Eduardo Marques-Carneiro¹, Rebeca Padrão Amorim¹, Michelle Gasparetti Leão Araujo¹, Astrid Nehlig²

¹Department of Neurology and Neurosurgery, Universidade Federal de Sao Paulo-UNIFESP, Rua Pedro de Toledo, 669 - 2º andar, CEP 04039-032, Sao Paulo, SP, Brazil, ²Inserm U-666, Strasbourg, France

TABLE OF CONTENTS

1. Abstract
2. Introduction
 - 2.1. Molecular changes in epilepsy
3. Adenosine: a brief review
 - 3.1. Adenosine and epilepsy
4. Erythropoietin
 - 4.1. Epo and epilepsy
5. Antiepileptic drugs
6. Conclusions
7. Acknowledgments
8. References

1. ABSTRACT

Temporal Lobe Epilepsy (TLE) is a chronic condition characterized by epileptic seizures originating mainly in temporal lobe areas. Epileptogenesis is a process in which a central nervous system injury can lead surviving neuronal populations to generate abnormal, synchronous and recurrent epileptiform discharges producing focal or generalized seizures. Hippocampal sclerosis, a massive cell death in the hippocampal formation and in the other regions of temporal lobe, is considered as hallmark of TLE. Despite the numerous antiepileptic drugs (AEDs) commercially available, about 30-40% of patients remain with seizures refractory to pharmacological treatment. In addition, there is no drug with significant efficacy to modify the epileptogenesis process. In this review we present some data regarding the neuroprotective effect of some adenosinergic agents, erythropoietin and carisbamate regarding the disease- and epileptogenesis-modifying effect.

2. INTRODUCTION

Epilepsy is a disorder of the brain characterized by an enduring predisposition to generate epileptic seizures and by the neurobiological, cognitive, psychological, and

social consequences of this condition (1). The term “epileptogenesis” refers to an injury-initiated change that causes surviving neuronal populations to generate abnormal, synchronous and recurring epileptiform discharges that produce focal or generalized behavioral seizures (2). However, Sloviter and Bumanglag (2) suggested that the window to act on the epileptogenesis process encompasses only the first week after *status epilepticus* (SE), and that the first days of the latent period correspond to a period during which a maturation of the epileptic circuitry occurs. Temporal Lobe Epilepsy (TLE) is a term used to define a chronic condition characterized by epileptic seizures originating preferentially, but not exclusively, in temporal lobe areas.

Neuropathological studies indicate that TLE is frequently associated with hippocampal sclerosis (HS) that is routinely detected by imaging studies during the presurgical evaluation of patients with this disorder (3). The histological pattern of HS includes loss of pyramidal cells in the prosubiculum, CA1, CA3 and hilus of dentate gyrus from the hippocampal formation (3). In addition, phenomena of synaptic rearrangement (sprouting of mossy fibers) and dispersion of

granule cells of the dentate gyrus are frequently observed in the HS from patients with TLE (4). Changes can also be found in other regions of the mesial temporal lobe, the entorhinal cortex and white matter (5). Patients who develop TLE demonstrate a progression in both the number of seizures and in the neurological symptoms related to the seizures, such as cognitive and behavioral disorders (6,7). The high prevalence and refractoriness to pharmacological treatment make this disease a subject of great interest for researchers in basic and clinical areas (8).

2.1. Molecular changes in epilepsy

Despite technological advances applied to neurosciences, little is known about the cellular and molecular phenomena related to epileptogenesis, the process by which a previously asymptomatic brain becomes capable of generating spontaneous seizures.

Several changes have been associated with epileptogenesis in hippocampus from patients with TLE or animal models of TLE. Hyper-excitability occurs in response to abnormal expression of GABAergic neurons in CA1 which can cause abnormal synchrony leading to seizure appearance (9). Reports of high expression of GluR1 in mossy cells of the hilus and in pyramidal neurons in CA3 have been associated with excitation of granule cells in hippocampus of patients with TLE (10).

Studies have reported that astrogliosis can also contribute to hyper-excitability and cell death resulting from seizures. Astrocytes contribute to the inflammatory response in the central nervous system (CNS). Glial cells can produce a variety of immunological molecules, such as class II major histocompatibility complex antigens, cytokines and chemokines. High levels of pro-inflammatory cytokines have been reported in brain areas involved with generation and propagation of seizures. Studies have shown that IL-1 β is up-regulated after 3h and 15 days in animals subjected to pilocarpine (11,12). This up-regulation is also observed in the sclerotic hippocampus of patients with TLE and can contribute to seizure onset through glutamate release (13,14). The genes regulated by IL-1 β are also up-regulated in the sclerotic hippocampus from TLE patients (15). IL-1 β is pro-ictogenic and the exposure to IL-1 β or TNF- α exacerbates the excitotoxic neuronal damage produced by NMDA and AMPA receptors

suggesting the role of cytokines in cell death. IL-1 β increases the calcium influx mediated by NMDA receptors thereby promoting excitotoxicity (16).

Despite the knowledge of a large number of mechanisms involved in hyper-excitability, seizure onset and cell damage resulting from seizures, the literature still lacks specific experimental models to study antiepileptogenesis. These models should allow the identification of brain areas involved with seizure triggering and what factors are responsible for this change.

To date, there is no report about any antiepileptic drug (AED) effective in treating seizures in patients with TLE which has been efficacious in preventing epileptogenesis (17). In this way, some strategies of neuroprotection with disease-modifying properties have been encouraged. Summarized below there is a brief review of studies on adenosinergic agents, erythropoietin and carisbamate regarding their neuroprotective and epilepsy-modifying action.

3. ADENOSINE: A BRIEF REVIEW

Adenosine is an endogenous nucleoside with modulatory action in different physiological processes in the CNS and in peripheral organs with potent neuroprotective action (18-21).

The regulation of the levels of intracellular and extracellular adenosine in the CNS occurs by the balance between the mechanisms of synthesis, release, uptake and degradation (21,22). In physiological conditions, the basal concentration of adenosine is relatively low in intracellular levels (10-50 nM) (23) and extracellular (30-300 nM) (21,24). However, its extracellular levels increase dramatically in conditions of metabolic stress such as ischemia, seizures or trauma (25).

The actions of adenosine in the CNS are mediated via P1 purinergic receptors, a family of G-protein coupled receptors. To date, four subtypes have been cloned and characterized: A1, A2A, A2B and A3 which differ by the type and the G-protein signal transduction triggering (21,24,25). Generally, the A1 and A3 receptors interact with inhibitory G proteins (Gi) promoting the inhibition of adenylate cyclase and thus decreasing the levels of cAMP. A2A and A2B receptors are coupled to stimulatory G protein (Gs) and activate adenylate cyclase, increasing cAMP (24).

Adenosine exhibits high affinity for receptors A1, A2A, A3 and low affinity for A2B receptor. The A1 and A2A receptors are most abundant in the CNS, where they are involved in most physiological activities. A2B and A3 receptors have lower expression in the CNS and may be relevant in some pathologic processes (26).

The A1 receptors are widely distributed in the CNS with pre- and post-synaptic location. They are highly expressed in the neocortex, hippocampus, cerebellum and spinal cord and display a low density in the striatum, amygdala, olfactory bulb, thalamus, substantia nigra and nucleus of the solitary tract (26). Although these receptors are most abundant in neurons, they are also present in astrocytes, microglia and oligodendrocytes (23).

The A2A receptors exhibit a more narrow distribution in the CNS compared to A1 receptors. They are predominantly expressed in the dorsal striatum, particularly in striato-pallidal GABAergic neurons and cortico-striatal glutamatergic terminals (23,25). However, these receptors are also found in other brain regions, including the neocortex, nucleus of the solitary tract, hippocampus and thalamus, albeit with lower levels of expression (26). As A1, A2A receptors are present in neurons, astrocytes, oligodendrocytes and microglia (23). They are also found in cerebral blood vessels (25). A2A receptors are present in the hippocampus, particularly in glutamatergic nerve terminals, favoring the release of glutamate (23). In presynaptic terminals A2A receptors may be co-localized with A1 receptors; thus, the activation of A2A reduces the inhibitory effects mediated by A1 (23, 27). Moreover, the presence of A2A receptors in the postsynaptic membrane of hippocampal neurons would promote depolarization of these cells (28).

Activation of A1 receptors is the most important inhibitory role played by adenosine in the CNS (25, 26). In contrast, A2A receptors are responsible for the excitatory actions of adenosine in the CNS and are involved in locomotion, anxiety, aggression, motivation and reinforcement in drug abuse and psychotic behaviors (29–31). Furthermore A2A receptors are involved in the control of cerebral blood flow (32).

3.1. Adenosine and epilepsy

In the CNS adenosine acts as a neuromodulator, predominantly inhibiting neuronal activity via A1 receptors (21,24). Thus, adenosine

has emerged as an important endogenous anticonvulsant agent (33). Microdialysis studies have reported that adenosine levels rise in the extracellular space of the hippocampus during the ictal phases in both experimental models and in patients with complex partial seizures and this increase has been proposed as an intrinsic mechanism for controlling seizures (33,34). Thus, any manipulation able to increase extracellular adenosine level offers significant potential for both preventing and blocking epileptic seizures (35).

In fact, the evidence suggests that the inhibitory effect of adenosine on seizure activity is mediated mainly by the activation of A1 receptors (18-20). These receptors have a high density in the hippocampus and are located in pre- and postsynaptic neuronal locations. Activation of presynaptic A1 receptors modulates excitatory synaptic transmission by decreasing glutamate release through inhibition of voltage-dependent calcium channels. In parallel, activation of postsynaptic A1 receptors depresses neuronal excitability by increasing membrane conductance to potassium ions hyperpolarizing the cell (21,35). These effects are crucial for the maintenance of intracellular calcium homeostasis and thus may protect nerve cells against excitotoxicity (25).

Several reports have shown that the adenosinergic A1 receptor agonist R-N6-phenylisopropyladenosine (R-PIA) and selective or non-selective A2A receptor antagonists are neuroprotective when administered prior to kainic acid (37) or pilocarpine (18,20,38,39). Accordingly, adenosinergic A1 receptor agonists can attenuate seizures in several experimental models of epilepsy as kindling (40–42), pilocarpine (43,44) or 3-nitropropionic acid (45), but the neuroprotective action was not reported by the authors. In fact, little is known about the action of these agents as modifiers of epileptogenesis.

Several studies have shown decreased density and activity of A1 receptors in epilepsy models (37–39). Since this inhibitory system is deficient it favors glutamate release and occurrence of seizures. Besides the decreased function of A1 receptors, reduction of the extracellular levels of adenosine due to the increase of adenosine kinase (ADK) activity in astrocytes has been reported in models of chronic epilepsy (49). Thus, strategies of activation of A1 receptors and manipulation of ADK in order to increase the extracellular level

of adenosine represent a challenge to control epileptogenesis.

Many authors have shifted their focus of interest to understanding the role of A2A receptor in modulating seizures (50). Contrary to what is observed with the A1 receptors, the density of A2A receptor is increased in the epileptic tissue (51). Furthermore, activation of the A2A receptor can stimulate microglia and astrocytes, causing the release of cytokines, increase of oxidative stress and inflammation (50), contributing to neurotoxicity and neuronal damage (27). Thus, blockade of A2A receptors has been considered a neuroprotective strategy for epilepsy (23,25,50).

According to some authors, the blockade of A2A receptors either using genetic deletion of A2A receptors in the pilocarpine model (52,53) or using non-selective antagonists such as caffeine administered chronically in the bicuculline and pentylentetrazol models (54), can cause a robust decrease in the severity of seizures. In contrast, the administration of the A2A antagonist, 3,7-dimethyl-1-propylxanthine (DMPX) prior to pilocarpine, increased the number of animals in SE (55). The chronic administration of caffeine prior to lithium-pilocarpine effectively prevented neuronal damage caused by seizures (38). However, the pretreatment with the selective A2A antagonist (7-(2-phenylethyl)-5-amino-2-(2-furyl)-pyrazolo-(4,3-e)-1,2,4-triazolo(1,5-c) pyrimidine) (SCH58261) did not change the pattern of seizure-induced cell death, although it reduced the number of animals presenting SE and the mortality, and increased the latency to SE onset (20).

Numerous studies targeted at understanding the role of A2A receptors in the modulation of epileptogenesis have generated controversial data highlighting the need of studies with new approaches.

Herein, we highlight the important role of adenosine as an endogenous inhibitor of neuronal excitability and neuroprotector, and we provide an overview about the role of each adenosinergic receptor in these process. Knowing the metabolic regulation of adenosine is the major point for understanding the modulation of adenosine on seizures or on epileptogenesis. Several authors detailed these points in reviews, especially Masino *et al.* (35). In summary, studies that show the role of adenosine in the modification of epileptogenesis are

still scarce in the literature although adenosine is emerging as a potent therapeutic target for epilepsy.

4. ERYTHROPOIETIN

Erythropoietin (EPO), a 34 kDa glycoprotein hormone, is produced primarily in the kidney and regulates the number of erythrocytes within the circulation to provide adequate tissue oxygenation (56-58). Recent studies have shown that EPO is a multifunctional molecule produced and utilized by many tissues. EPO is induced under hypoxia, hypoglycemia, strong neuronal depolarization and excess of oxygen radicals (59-61). The induction of the EPO gene in most tissues is regulated by the hypoxia-inducible factor-1 (HIF-1) (62,63). Both the EPO and EPO receptor (EPOR) can be expressed in various organs, as rodent and human brain, as well as in cultured neurons, astrocytes, oligodendrocytes, microglia, and endothelial cells, and they are related to endogenous neuroprotective effect (64-70).

EPO is present in the CNS and also in the vascular and immune systems. In these systems, EPO offers robust protection against cell death caused by oxidative stress, excitotoxicity and inflammation. In addition, EPO promotes neurogenesis and differentiation in the brain and induces angiogenesis via downstream effectors such as vascular endothelial growth factor (VEGF) (56,57,71-75).

EPOR is expressed constitutively in almost all neurons of the hippocampus, except within the hilus (63). Neuronal EPOR immunolabeling is concentrated within cell bodies and varicosities, except in the CA1 area where EPOR is also found in basal dendrites of pyramidal neurons laying throughout the *stratum radiatum*. The basal expression of EPOR in the hippocampus suggests that it plays a role in neuronal homeostasis (63).

The ability of peripherally administered EPO as well as recombinant human EPO (rhEPO) to cross the blood-brain barrier (BBB), combined with its robust neurotrophic and angiogenic efficacy, has sparked interest in EPO as a potential neuroprotector (61,67,69,70,73,76-78). The administration of exogenous EPO *in vivo* or *in vitro* preparations has shown neurotrophic and neuroprotective actions against central and peripheral neuronal injury associated with trauma, stroke, ischemia, inflammation and epileptic seizures (56,58,61,63,72,73,76,79-82).

4.1. EPO and epilepsy

In epilepsy models, EPO has been associated with a variety of functions including reduction of seizure duration, protection against BBB leakage and microglial activation, prevention of aberrant cell genesis and granule cell dispersion, reduction in infiltration of macrophages and in inflammatory cytokines, decreased neuronal damage associated with spontaneous recurrent seizures (SRS) in rats (56,58,61,63,65,70,72–74,78,83–85).

Considering the neuroprotective and anti-inflammatory potential of EPO, some authors tested EPO-derived mimetic peptides regarding its disease modifying or antiepileptogenic ability. When administrated after SE onset, the peptide pyroglutamate helix B surface peptide (pHBSP) promoted hippocampal cell proliferation, neuronal differentiation and cell survival but did not affect the number of rats presenting spontaneous seizures (86). However, pHBSP was able to attenuate mood disorders as well as cognitive deficits associated to epilepsy (86).

In this line, Epotris, another EPO-derived peptide devoid of erythropoietic activity, was administrated to evaluate the histopathological consequences of SE induced by electrical stimulation (69). Epotris attenuated the seizure-associated expansion of the neuronal progenitor cell population and affected the number of basal dendrites in these progenitor cells (69). In addition, Epotris diminished the microglial activation caused by seizures in the thalamus, but did not interfere with hippocampal cell loss. According to the authors, Epotris exerted limited *in vivo* effects on cell consequences caused by prolonged seizures (69). Despite the opposing biological effects, the EPO-derived peptide mimetic design offers intriguing possibilities as therapeutic strategy since it may yield molecules with disease- or epileptogenesis-modifying properties (87).

Several studies have shown enhancement of EPOR in the hippocampus following SE induced by pilocarpine, lithium-pilocarpine or kainic acid (56,60,63,65,78). When administered after SE onset, EPO provides neural protection in the CA1, CA3, and hilus (56,60,63,65,78). According to the authors, the mechanisms underlying this effect can be associated with caspase-3 inhibition, increase in the expression of Bcl-w, elevated expression of Bcl-XL, normal expression of Bim, and up-regulation of Bcl-2, which can neutralize Bid, a pro-apoptotic

Bcl-2 protein, and cause down-regulation of Bax (39,65,84,88,89). In addition, EPO is able to decrease glutamate release, increase GABA release and suppress the influx of calcium via PI3K/Akt and/or ERK1/2 signaling pathway, increasing the expression of p-Akt protein which in turn regulates the expression of caspase-9, hence protecting neurons against excitotoxicity (60,71,61,77,89). Given these data, we consider that treatment with EPO offers exciting opportunities to prevent the onset and progression of neurodegenerative disorders as epilepsy.

5. ANTIEPILEPTIC DRUGS

AEDs are the most common treatment of epilepsy (91). However, even with more than 10 new AEDs available for purchase, about 30-40% of patients with TLE remain with refractory seizures even in polytherapy (92). A variety of mechanisms of action has been assigned to new and conventional AEDs, including modulation of ion channels, GABAergic and glutamatergic neurotransmission (metabolism, receptor and secondary messengers) (93). In some cases, more than one mechanism can be attributed to one AED. For example, phenobarbital increases GABA levels, potentiates GABA_A receptors and facilitates chloride ion flux (91). Several authors have focused their interest on studying the modulation of epilepsy by AEDs and some data are shown in Table 1 (93).

As can be seen in Table 1, there is a considerable literature showing that AEDs might have a beneficial effect in the prevention of neuronal death resulting from epileptic seizures (94). Different patterns of neuroprotection can be seen with one AED depending on the epilepsy model tested (Table-1). For example diazepam applied in the pilocarpine model induced neuroprotection in CA1, CA3 and hilus (95). However in the kainic acid model, the neuroprotection induced by diazepam was observed in the amygdala, piriform cortex and endopiriform nucleus (96). In the kindling model diazepam induced a significant neuroprotection in CA1, CA3 and hilus only when administrated 2h after SE (97) (Table-1). Besides neuroprotection, diazepam induced significant SRS changes in a few cases, modifying the duration, frequency and latency of seizures (97–104).

The AED carisbamate (RWJ-333369; (S)-2-O-carbamoyl-1-ochlorophenyl-ethanol), has received special attention due to its ability

Table 1. Neuroprotective AEDs and their effect on seizures and epileptogenesis

AED	Model	Beginning of treatment	Duration of treatment	Doses Mg/kg	Effect on epileptogenesis	Effect on seizures	Neuroprotection	Ref.
DZP	Pilocarpine	4, 28, 52 and 76 h after SE	Unique	2	ND	ND	CA1, CA3 and hilus	(95)
	Kainic acid	3h after SE	Unique	25	ND	ND	amygdala, piriform cortex and endopiriform nucleus	(114)
	Amygdala kindling	2 or 3h after SE	Unique	20	Only 42% of rats developed SRS	↓ frequency	CA1, CA3 and hilus only in 2h after SE	(97)
VPA	SSSE	4h after SE (SE was stopped with diazepam)	4 weeks	400-200 (3 × day)	No change	↓ frequency	hippocampus	(98)
CBZ	Pilocarpine	ND	56 days	40 (3 × day)	No change	↓ frequency and duration	hippocampus	(99)
	Pilocarpine	4, 28, 52 and 76 h after SE	Unique	120	ND	ND	CA1, CA3 and hilus	(95)
PHT	Pilocarpine	4, 28, 52 and 76 h after SE	Unique	60	ND	ND	CA1, CA3 and hilus	(95)
FBM	PPS	10 and 40 min after PPS	Unique	50, 100 and 200	ND	ND	CA1, piriform cortex, subiculum and amygdala	(115)
LTG	PPS	1h after PPS	2 weeks	12.5. (2 × day)	ND	ND	CA3, hilus and piriform cortex	(116)
TPM	Kindling (HPC)	140 min after stimulation	Unique	20, 40 and 80	ND	ND	CA1, hilus and CA3 (contralateral)	(114)
	Lithium-Pilocarpine	10h after SE	6 days	10, 30 and 60 (2 × day)	No change	No change	CA1 and CA3 but not hilus, entorhinal and piriform cortex	(117)
VGB	Lithium-Pilocarpine	10 min after Pilocarpine administration	45 days	250	No change	No change	CA3, CA1 and hilus	(118)
PGB	Lithium-Pilocarpine	20 min after pilocarpine	7 days	50	No change	increased latency for SRS	Piriform and entorhinal cortex	(102)
TGB	PPS	Immediately after evoked potentials	4 days	50	ND	ND	CA1 and CA3	(119)
LSM	SSSE	24h after SE (Stopped with diazepam)	24 days	10 and 30 (3 × day)	No changes	No change	CA1 and piriform cortex only at the dose of 30 mg/kg	(120)
CRS	Lithium-Pilocarpine	1h after SE	7 days	30, 60, 90 and 120 (2 × day)	Suppressed SRS 50% of rats	Delayed the SRS on the other 50% of rats	CA1, CA3, thalamus, amygdala, entorhinal and piriform cortex	(104)
	Lithium-Pilocarpine	1h after SE	7 days	90 (2 × day)	Suppressed SRS 50% of rats		CA1, CA3, thalamus, amygdala, entorhinal and piriform cortex	(103)

Perforant-path stimulation (PPS); self-sustained *status epilepticus* (SSSE); spontaneous recurrent seizures (SRS); amygdala (AMY); hippocampus (HPC); diazepam (DZP); valproate (VPA); carbamazepine (CBZ); phenobarbital (PHB); phenytoin (PHT); felbamate (FBM); lamotrigine (LTG); topiramate (TPM); vigabatrin (VGB); pregabalin (PGB); tiagabine (TGB); lacosamide (LSM); carisbamate (CRB)

to modify the epileptogenesis besides to induce significant neuroprotection. Indeed, carisbamate is an anticonvulsant with a wide spectrum of activity and a substantial safety margin (105). However, little is known about its mechanism of action. Some authors have reported that carisbamate blocks the voltage-gated sodium channel (106), although other authors demonstrated that carisbamate inhibits excitatory synaptic strength by a presynaptic mechanism without affecting the GABAergic system (107). In a recent study, Shin *et al.* (108) reported that treatments during 2 or 14 days with carisbamate decrease the firing activity in the dorsal raphe nucleus, locus coeruleus and ventral tegmental area, probably due to a reduction in the serotonin (5HT), norepinephrine and dopamine neurotransmission. Nevertheless, in the same study the authors reported increased activation of 5-HT_{1A} receptors in the hippocampus (108).

In post-SE epilepsy models such as kainic acid and lithium-pilocarpine, carisbamate presents an anticonvulsant action by significantly increasing the latency to the occurrence of SRS (109,110). In the lithium-pilocarpine model, SRS occurrence was observed only in 50% of carisbamate treated rats (110). Other authors using the lithium-pilocarpine model have shown that the administration of carisbamate (90 and 120 mg/kg, i.p.), at 1 and 9 hours after SE onset, over 6 days, produces a strong neuroprotection in the hippocampus, entorhinal and piriform cortices, thalamus, amygdala, nucleus basalis magnocellularis as well in orbital, infralimbic and prelimbic cortices (103,104,111). In addition a significant preservation of learning and memory, attention, locomotion and coordination capabilities was recorded when animals were subjected to behavioral testing (103,111). In this model, carisbamate prevented mossy fiber sprouting in the dentate gyrus and suppressed SRS in almost 50% of treated rats (104). Interestingly, rats that did not present characteristic SRS displayed spike-and-wave discharges, an electrographic pattern characteristic of absence seizures. In addition, carisbamate exhibited anti-seizure effect in the Genetic Absence Epilepsy Rat from Strasbourg (GAERS) since it abolished spike-and-wave discharges (112). These data show the potential disease-modifying effect of carisbamate

Based on these data, carisbamate has been considered a strong epilepsy-modifying AED with neuroprotective properties whose mechanisms

need to be elucidated. To date, several studies are ongoing to attempt understanding more about the cellular and molecular mechanisms underlying carisbamate action. The proteomic technology using 2-dimensional electrophoresis to determine differential protein expression in brain areas of the epileptic circuit of rats treated with carisbamate have shown changes in glycolytic pathways associated with carisbamate treatment. Besides, differential activation of thalamic nuclei involved with seizure spread, has been observed in carisbamate-treated rats compared to untreated rats. Finally, partial results have shown that carisbamate also alters the level of monoamines (serotonin, dopamine and noradrenaline) and its metabolites in the hippocampus and entorhinal/piriform cortices of rats (data in progress).

In summary, despite numerous AEDs commercially available, about 30-40% of patients with TLE remain with seizures refractory to pharmacological treatment (92). New AEDs have been developed in an attempt to modify SRS occurrence but also to reduce the side effects of classical AEDs (113). Considerable neuroprotection has been observed with the use of AEDs in experimental models of epilepsy, however, little effect on the expression of SRS or epileptogenesis is reported (see Table 1). Interestingly, carisbamate presents neuroprotective effect in post-SE model of epilepsy, exerts a disease-modifying effect, since it increases the latency to SRS onset and presents an epileptogenic-modifying effect, since it may induce spike-and-wave discharges (SWD) instead of limbic seizures in the pilocarpine model. In addition, carisbamate induces a significant behavior and cognitive preservation in the pilocarpine model. Altogether, carisbamate can be considered as very promising tool to study epileptogenic-modifying mechanisms in epilepsy.

6. CONCLUSIONS

The development of new strategies able to promote neuroprotection and to modify the epileptogenesis is a topic of high interest. Strategies which allow administrating some molecules in the earlier stage of the epileptogenic process of a post-SE model, and then evaluate the impact on SRS latency or frequency, are promising strategies to find potential therapeutic targets. In this way, adenosinergic agents, EPO and carisbamate are promising molecules, and can bring new knowledge

on mechanisms underlying epileptogenesis.
<http://www.sciencedirect.com/science/article/pii/S1059131111000124> - bib0230.

7. ACKNOWLEDGMENTS

The authors thank to Fapesp, INNT/MCT/CNPq, CAPES for financial support.

8. REFERENCES

1. R.S. Fisher, W.V.E. Boas, W. Blume, C. Elger, P. Genton, P. Lee, J. Engel: Epileptic Seizures and Epilepsy: Definitions Proposed by the International League Against Epilepsy (ILAE) and the International Bureau for Epilepsy (IBE). *Epilepsia* 46, 470-472 (2005)
DOI: 10.1111/j.0013-9580.2005.66104.x
2. R.S. Sloviter, A.V. Bumanglag: Defining "epileptogenesis" and identifying "antiepileptogenic targets" in animal models of acquired temporal lobe epilepsy is not as simple as it might seem. *Neuropharmacology* 69, 3-15 (2013)
DOI: 10.1016/j.neuropharm.2012.01.022
3. G.W. Mathern, J.K. Pretorius, T.L. Babb: Influence of the type of initial precipitating injury and at what age it occurs on course and outcome in patients with temporal lobe seizures. *J Neurosurg* 82(2), 220-227 (1995)
DOI: 10.3171/jns.1995.82.2.0220
4. T.L. Babb, W.R. Kupfer, J.K. Pretorius, P.H. Crandall, M.F. Levesque: Synaptic reorganization by mossy fibers in human epileptic fascia dentata. *Neuroscience* 42(2), 351-363 (1991)
DOI: 10.1016/0306-4522(91)90380-7
5. B.S. Kasper, H. Stefan, M. Buchfelder, W. Paulus: Temporal lobe microdysgenesis in epilepsy versus control brains. *J Neuropathol Exp Neurol* 58(1), 22-28 (1999)doi :10.1097/00005072-199901000-00003
6. J. Jr. Engel, R. Bandler, N.C. Griffith, S. Caldecott-Hazard: Neurobiological evidence for epilepsy-induced interictal disturbances. *Adv Neurol* 55, 97-111 (1991)
doi not found
7. J.A. French, A.M. Kanner, J. Bautista, B. Abou-Khalil, T. Browne, C.L. Harden: Efficacy and tolerability of the new antiepileptic drugs I: treatment of new onset epilepsy. *Neurology* 62, 1252-1260 (2004)
DOI:10.1212/01.WNL.0000123693.82339.FC
8. L.M. Li, J.W. Sander: National demonstration project on epilepsy in Brazil. *Arq Neuropsiquiatr* 61(1), 153-156 (2003)
DOI:10.1590/S0004-282X2003000100033
9. L. Wittner, L. Eross, Z. Szabó, S. Tóth, S. Czirják, P. Halász, T.F. Freund, Z.S. Maglóczy: Synaptic reorganization of calbindin-positive neurons in the human hippocampal CA1 region in temporal lobe epilepsy. *Neuroscience* 115(3), 961-978 (2002)
DOI: 10.1016/S0306-4522(02)00264-6
10. T. Eid, I. Kovacs, D.D. Spencer, N.C. de Lanerolle: Novel expression of AMPA-receptor subunit GluR1 on mossy cells and CA3 pyramidal neurons in the human epileptogenic hippocampus. *Eur J Neurosci* 15(3), 517-527 (2002)
DOI: 10.1046/j.0953-816x.2001.01887.x
11. T.L.F. Gouveia, F.A. Scorza, M.J.V. Silva, T.A. Bandeira, S.R. Perosa, G.A. Argañaraz, M. P. Silva, T.R. Araujo, M.L. Frangiotti, D. Amado, E.A. Cavalheiro, J.A.Jr. Silva, M. G. Naffah-Mazzacoratti: Lovastatin decreases the synthesis of inflammatory mediators in the hippocampus and blocks the hyperthermia of rats submitted to long-lasting status epilepticus. *Epilepsy Behav* 20(1), 1-5 (2011)
DOI: 10.1016/j.yebeh.2010.10.001
12. T.L.F. Gouveia, F.A. Scorza, H.A. Iha, M.I.B. Frangiotti, S.R. Perosa, E.A. Cavalheiro, J.A.Jr. Silva, R.S. Feliciano,

- A.C. Almeida, M.G. Naffah-Mazzacoratti: Lovastatine decreases the synthesis of inflammatory mediators during epileptogenesis in hippocampus of rat submitted to pilocarpine-induced epilepsy. *Epilepsy Behav* (In Press)
DOI not found
13. G.R. John, S.C. Lee, X. Song, M. Rivieccio, C.F. Brosnan: IL-1-regulated responses in astrocytes: relevance to injury and recovery. *Glia* 49(2), 161-176 (2005)
DOI: 10.1002/glia.20109
14. P.P.V. Varella, J.F.C. Santiago, H.Jr. Carrete, E.M.S. Higa, E.M.T. Yacubian, R.S. Centeno, L.O. Caboclo, E.F. Castro Neto, M. Canzian, D. Amado, E.A. Cavalheiro, M. G. Naffah-Mazzacoratti: Relationship between fluid-attenuated inversion-recovery (FLAIR) signal intensity and inflammatory mediator's levels in the hippocampus of patients with temporal lobe epilepsy and mesial temporal sclerosis. *Arq Neuropsiquiatr* 69(1),91-99 (2011)
DOI: 10.1590/S0004-282X2011000100018
15. T. Ravizza, B. Gagliardi, F. Noé, K. Boer, E. Aronica, A. Vezzani: Innate and adaptive immunity during epileptogenesis and spontaneous seizures: evidence from experimental models and human temporal lobe epilepsy. *Neurobiol Dis* 29(1), 142-160 (2007)
DOI: 10.1016/j.nbd.2007.08.012
16. A. Vezzani, S. Balosso, T. Ravizza: The role of cytokines in the pathophysiology of epilepsy. *Brain Behav Immun* 22(6), 797-803 (2008)
DOI: 10.1016/j.bbi.2008.03.009
17. N.R. Temkin: Preventing and treating posttraumatic seizures: the human experience. *Epilepsia* 50(2), 10-13 (2009)
DOI: 10.1111/j.1528-1167.2008.02005.x
18. I.R. Silva, A. Nehlig, F.E. Rosim, T. Vignoli, D.S. Persike, J.P. Blini, E.A. Cavalheiro: Epilepsia e neuroproteção: o papel do agonista adenosinérgico A1 (R-Pia) na modulação da crise induzida por pilocarpina. *J Epilepsy Clin Neurophysiol* 14, 106-110 (2008)
DOI: 10.1590/S1676-26492008000300005
19. I.R. Silva, A. Nehlig, F.E. Rosim, T. Vignoli, D.S. Persike, A. Ferrandon, R. Sinigaglia-Coimbra, M.J.S. Fernandes: The A1 receptor agonist R-Pia reduces the imbalance between cerebral glucose metabolism and blood flow during status epilepticus: Could this mechanism be involved with neuroprotection? *Neurobiol Dis* 41, 169-176 (2011)
DOI: 10.1016/j.nbd.2010.09.004
20. F.E. Rosim, D.S. Persike, A. Nehlig, R.P. Amorim, D.M. Oliveira, M.J.S. Fernandes: Differential neuroprotection by A1 receptor activation and A2A receptor inhibition following pilocarpine-induced status epilepticus. *Epilepsy Behav* 22, 207-13 (2011)
DOI: 10.1016/j.yebeh.2011.07.004
21. T.V. Dunwiddie, S.A. Masino: The role and regulation of adenosine in the central nervous system. *Annu Rev Neurosci* 24, 31-55 (2001)
DOI: 10.1146/annurev.neuro.24.1.31
22. S. Latini, F. Pedata: Adenosine in the central nervous system: release mechanisms and extracellular concentrations. *J Neurochem* 79, 463-484 (2001)
DOI: 10.1046/j.1471-4159.2001.00607.x
23. R.A. Cunha: Neuroprotection by adenosine in the brain: from A1 receptor activation to A2A receptor blockade. *Purinergic Signal* 1, 111-134 (2005)
DOI: 10.1007/s11302-005-0649-1
24. B.B. Fredholm, K. Bättig, J. Holmén, A. Nehlig, E.E. Zvartau: Actions of caffeine in the brain with special reference to factors that contribute to its widespread use. *Pharmacol Rev* 51, 83-133 (1999)
doi not found
25. E.E. Benarroch: Adenosine and its receptors. Multiple modulatory functions and potential therapeutic targets for neurologic disease. *Neurology* 70,

- 231-236 (2008)
DOI: 10.1212/01.wnl.0000297939.18236.ec
26. J.A. Ribeiro, A.M. Sebastião, A. Mendonça: Adenosine receptors in the nervous system: pathological implications. *Prog Neurobiol* 68, 377-392 (2003)
DOI: 10.1016/S0301-0082(02)00155-7
27. De Mendonça, A.M. Sebastião, J.A. Ribeiro: Adenosine: does it have a neuroprotective role after all? *Brain Res Brain Rev* 33, 258-274 (2000)
DOI: 10.1016/S0165-0173(00)00033-3
28. H. Li, J.L. Henry: Adenosine A2A receptor mediation of pre- and postsynaptic excitatory effects of adenosine in rat hippocampus *in vitro*. *Eur J Pharmacol* 347, 173-182 (1998)
DOI: 10.1016/S0014-2999(98)00105-8
29. C. Ledent, J.M. Vaugeois, S.N. Schiffmann, T. Pedrazzini, M. El Yacoubi, J.J. Vanderhaeghen, J. Costentin, J.K. Heath, G. Vassart, M. Parmentier: Agressiveness, hypoalgesia and high blood pressure in mice lacking the adenosine A2A receptor. *Nature* 388, 674-678 (1997)
DOI: 10.1038/41771
30. J.L. Moreau, G. Huber: Central adenosine A2A receptors: an overview. *Brain Res* 31, 65-82 (1999)
DOI: 10.1016/S0165-0173(99)00059-4
31. J.F. Chen, M. Beilstein, Y.H. Xu, T.J. Turner, R. Moratalla, D.G. Standaert, V.J. Aloyo, J.S. Fink, M.A. Schwarzschild: Selective attenuation of psychostimulant-induced behavioral responses in mice lacking A2A adenosine receptors. *Neuroscience* 97, 195-204 (2000)
DOI: 10.1016/S0306-4522(99)00604-1
32. V. Ralevic, G. Burnstock: Receptors for purines and pyrimidines. *Pharmacol Rev* 50, 413-492 (1998)
doi not found
33. O. Pagonopoulou, A. Efthimiadou, B. Asimakopoulos, N.K. Nikolettos: Modulatory role of adenosine and its receptors in epilepsy: possible therapeutic approaches. *Neurosci Res* 56, 14-20 (2006)
DOI: 10.1016/j.neures.2006.05.010
34. M.J. During, D.D. Spencer: Adenosine: a potential mediator of seizure arrest and postictal refractoriness. *Ann Neurol* 32, 618-624 (1992)
DOI: 10.1002/ana.410320504
35. S.A. Masino, M. Kawamura, C.D. Wasser, C.A. Wasser, L.T. Pomeroy, D.N. Ruskin: Adenosine, ketogenic diet and epilepsy: the emerging therapeutic relationship between metabolism and brain activity. *Curr Neuropharmacol* 7(3), 257-268 (2009)
DOI: 10.2174/157015909789152164
36. E. Ongini, P. Schubert: Neuroprotection induced by stimulating A1 or blocking A2A adenosine receptors: an apparent paradox. *Drug Dev Res* 45, 387-393 (1998)
DOI: 10.1002/(SICI)1098-2299(199811/12)45:3/4<387::AID-DDR39>3.0.CO;2-U
37. D.G. MacGregor, T.W. Stone: Inhibition by the adenosine analogue, (R)-N6-phenylisopropyladenosine, of kainic acid neurotoxicity in rat hippocampus after systemic administration. *Br J Pharmacol* 109(2), 316-321 (1993)
DOI: 10.1111/j.1476-5381.1993.tb13572.x
38. M.A. Rigoulot, C. Leroy, E. Koning, A. Ferrandon, A. Nehlig: Prolonged Low-dose Caffeine Exposure Protects Against Hippocampal Damage but Not Against the Occurrence of Epilepsy in the Lithium-pilocarpine Model in the Rat. *Epilepsia* 44(4), 529-535 (2003)
DOI: 10.1046/j.1528-1157.2003.50502.x
39. S. Bartesaghi, M. Marinovich, E. Corsini, C.L. Galli, B. Viviani: Erythropoietin: a novel neuroprotective cytokine. *Neurotoxicology* 26, 923-928 (2005)
DOI: 10.1016/j.neuro.2005.01.016
40. M. Pourgholami, J. Mirnajafi-Zadeh, J. Behzadi: Effect of intraperitoneal and

- intrahippocampal (CA1) 2-chloroadenosine in amygdaloid kindled rats. *Brain Res* 751(2), 259-264 (1997)
DOI: 10.1016/S0006-8993(96)01406-0
41. A.S. Abdul-Ghani, P.J.E. Attwell, H.F. Bradford: The protective effect of 2-chloroadenosine against the development of amygdala kindling and on amygdala-kindled seizures. *Eur J Pharmacol* 326(1), 7-14 (1997)
DOI: 10.1016/S0014-2999(97)00139-8
42. M. Zeraati, J. Mirnajafi-Zadeh, Y. Fathollahi, S. Namvar, M.E. Rezvani: Adenosine A1 and A2A receptors of hippocampal CA1 region have opposite effects on piriform cortex kindled seizures in rats. *Seizure* 15(1), 41-48 (2006)
DOI: 10.1016/j.seizure.2005.10.006
43. W.A. Turski, E.A. Cavalheiro, C. Ikonomidou, L.E. Mello, Z.A. Bortolotto, L. Turski: Effects of aminophylline and 2-chloroadenosine on seizures produced by pilocarpine in rats: morphological and electroencephalographic correlates. *Brain Res* 361(1-2), 309-323 (1985)
DOI: 10.1016/0006-8993(85)91302-2
44. G.M. Khan, I. Smolders, G. Ebinger, Y. Michotte: Anticonvulsant effect and neurotransmitter modulation of focal and systemic 2-chloroadenosine against the development of pilocarpine-induced seizures. *Neuropharmacology* 39(12), 2418-2432 (2000)
DOI: 10.1016/S0028-3908(00)00072-1
45. B. Zuchora, M. Wielosz, E.M. Urbańska: Adenosine A1 receptors and the anticonvulsant potential of drugs effective in the model of 3-nitropropionic acid-induced seizures in mice. *Eur Neuropsychopharmacol* 15(1), 85-93 (2005)
DOI: 10.1016/j.euroneuro.2004.05.006
46. M. Glass, R.L. Faull, J.Y. Bullock, K. Jansen, E.W. Mee, E.B. Walker, B.J. Synek, M. Dragunow: Loss of A1 adenosine receptors in human temporal lobe epilepsy. *Brain Res* 710, 56-68 (1996)
DOI: 10.1016/0006-8993(95)01313-X
47. T. Ochiishi, M. Takita, M. Ikemoto, H. Nakata, S.S. Suzuki: Immunohistochemical analysis on the role of adenosine A1 receptors in epilepsy. *NeuroReport* 10, 3535-3541 (1999)
DOI: 10.1097/00001756-199911260-00014
48. A. Ekonomou, G. Sperk, G. Kostopoulos, F. Angelatos: Reduction of A1 adenosine receptors in rat hippocampus after kainic acid-induced limbic seizures. *Neurosci Lett* 284, 49-52 (2000)
DOI: 10.1016/S0304-3940(00)00954-X
49. D. Boison: The adenosine kinase hypothesis of epileptogenesis. *Prog Neurobiol* 84(3), 249-262 (2008)
DOI: 10.1016/j.pneurobio.2007.12.002
50. J. Wardas: Adenosine and Neuroprotection. *Pol J Pharmacol* 54, 313-326 (2002)
DOI not found
51. N. Rebola, R.J. Rodrigues, L.V. Lopes, P.J. Richardson, C.R. Oliveira, R.A. Cunha: Adenosine A1 and A2A receptors are co-expressed in pyramidal neurons and co-localized in glutamatergic nerve terminals of the rat hippocampus. *Neuroscience* 133(1), 79-83 (2005)
DOI: 10.1016/j.neuroscience.2005.01.054
52. M. El Yacoubi, C. Ledent, M. Parmentier, J. Costentin, J.M. Vaugeois: Adenosine A2A receptor deficient mice are partially resistant to limbic seizures. *Naunyn Schmiedeberg's Arch Pharmacol* 380(3), 223-232 (2009)
DOI: 10.1007/s00210-009-0426-8
53. M. El Yacoubi, C. Ledent, M. Parmentier, J. Costentin, J.M. Vaugeois: Evidence for the involvement of the adenosine A(2A) receptor in the lowered susceptibility to pentylenetetrazol-induced seizures produced in mice by long-term treatment with caffeine. *Neuropharmacology* 55(1), 35-40 (2008)
DOI: 10.1016/j.neuropharm.2008.04.007
54. B. Johansson, V. Georgiev, T. Kuosmanen, B.B. Fredholm: Long-term treatment

- with some methylxanthines decreases the susceptibility to bicuculline- and pentylenetetrazol-induced seizures in mice. Relationship to c-fos expression and receptor binding. *Eur J Neurosci* 8(12), 2447-2458 (1996)
DOI: 10.1111/j.1460-9568.1996.tb01539.x
55. E.P.M. Vianna, A.T. Ferreira, F. Doná, E.A. Cavalheiro, M.J.S. Fernandes: Modulation of seizures and synaptic plasticity by adenosinergic receptors in an experimental model of temporal lobe epilepsy induced by pilocarpine in rats. *Epilepsia* 46(5), 166-173 (2005)
DOI: 10.1111/j.1528-1167.2005.01027.x
 56. K. Chu, K.H. Jung, S.T. Lee, J.H. Kim, K.M. Kang, H.K. Kim, J.S. Lim, H.K. Park, M. Kim, S.K. Lee, J.K. Roh: Erythropoietin reduces epileptogenic processes following status epilepticus. *Epilepsia* 49, 1723-1732 (2008)
DOI: 10.1111/j.1528-1167.2008.01644.x
 57. K. Maiese, F. Li, Z.Z. Chong: Erythropoietin in the brain: can the promise to protect be fulfilled? *Trends Pharmacol Sci* 25, 577-583 (2004)
DOI: 10.1016/j.tips.2004.09.006
 58. M. Jing, T. Shingo, T. Yasuhara, A. Kondo, T. Morimoto, F. Wang, T. Baba, W.J. Yuan, N. Tajiri, T. Uozumi, M. Murakami, M. Tanabe, Y. Miyoshi, S. Zhao, I. Date: The combined therapy of intrahippocampal transplantation of adult neural stem cells and intraventricular erythropoietin-infusion ameliorates spontaneous recurrent seizures by suppression of abnormal mossy fiber sprouting. *Brain Res* 1295, 203-217 (2009)
DOI: 10.1016/j.brainres.2009.07.079
 59. M. Chikuma, S. Masuda, T. Kobayashi, M. Nagao, R. Sasaki: Tissue-specific regulation of erythropoietin production in the murine kidney, brain and uterus. *Am J Physiol - Endocrinol Metab* 279, 1242-1248 (2000)
doi not found
 60. P.E. Sanchez, R.P. Fares, J.J. Risso, C. Bonnet, S. Bouvard, M. Le-Cavorsin, B. Georges, C. Moulin, A. Belmeguenai, J. Bodennec, A. Morales, J.M. Pequignot, E.E. Baulieu, R.A. Levine, L. Bezin: Optimal neuroprotection by erythropoietin requires elevated expression of its receptor in neurons. *Proc Natl Acad Sci USA* 106, 9848-9853 (2009)
DOI: 10.1073/pnas.0901840106
 61. G. Üzümlü, A. Sarper Diler, N. Bahçekapili, Y. Ziya Ziyil: Erythropoietin prevents the increase in blood-brain barrier permeability during pentylenetetrazol induced seizures. *Life Sci* 78, 2571-2576 (2006)
DOI: 10.1016/j.lfs.2005.10.027
 62. H.H. Marti: Erythropoietin and the hypoxic brain. *J Exp Biol* 207, 3233-3242 (2004)
DOI: 10.1242/jeb.01049
 63. J. Nadam, F. Navarro, P. Sanchez, C. Moulin, B. Georges, A. Laglaine, J.M. Pequignot, A. Morales, P. Ryvlin, L. Bezin: Neuroprotective effects of erythropoietin in the rat hippocampus after pilocarpine-induced status epilepticus. *Neurobiol Dis* 25, 412-426 (2007)
DOI: 10.1016/j.nbd.2006.10.009
 64. K. Chin, X. Yu, B. Beleslin-Cokic, C. Liu, K. Shen, H.W. Mohrenweiser, C.T. Noguchi: Production and processing of erythropoietin receptor transcripts in brain. *Brain Res Mol Brain Res* 81, 29-42 (2000)
DOI: 10.1016/S0169-328X(00)00157-1
 65. Kondo, T. Shingo, T. Yasuhara, S. Kuramoto, M. Kameda, Y. Kikuchi, T. Matsui, Y. Miyoshi, T. Agari, C.V. Borlongan, I. Date: Erythropoietin exerts anti-epileptic effects with the suppression of aberrant new cell formation in the dentate gyrus and upregulation of neuropeptide Y in seizure model of rats. *Brain Res* 1296, 127-136 (2009)
DOI: 10.1016/j.brainres.2009.08.025
 66. E. Morishita, S. Masuda, M. Nagao, Y. Yasuda, R. Sasaki: Erythropoietin receptor is expressed in rat hippocampal and cerebral cortical neurons, and

- erythropoietin prevents *in vitro* glutamate-induced neuronal death. *Neuroscience* 76, 105-116 (1997)
DOI: 10.1016/S0306-4522(96)00306-5
67. C.T. Noguchi, P. Asavaritikrai, R. Teng, Y. Jia: Role of erythropoietin in the brain. *Crit Rev Oncol Hematol* 64, 159-171 (2007)
DOI: 10.1016/j.critrevonc.2007.03.001
68. A.L. Sirén, F. Knerlich, W. Poser, C.H. Gleiter, W. Bruck, H. Ehrenreich: Erythropoietin and erythropoietin receptor in human ischemic/hypoxic brain. *Acta Neuropathol (Berl)* 101, 271-276 (2001)
DOI not found
69. C. Zellinger, N. Seeger, M. Hadamitzky, S. Fischborn, V. Russmann, H. Wendt, S. Pankratova, E. Bock, V. Berezin, H. Potschka: Impact of the erythropoietin-derived peptide mimetic Epotris on the histopathological consequences of status epilepticus. *Epilepsy Res* 96, 241-249 (2011)
DOI: 10.1016/j.epilepsyres.2011.06.009
70. H. Zheng, X. Wang, Z. Tang, W. Zheng, Z. Li: The PI3K/Akt and ERK1/2 signaling pathways mediate the erythropoietin-modulated calcium influx in kainic acid-induced epilepsy. *Neuroreport* 24, 335-341 (2013)
DOI: 10.1097/WNR.0b013e32835ffe03
71. S. Chateauvieux, C. Grigorakaki, F. Morceau, M. Dicato, M. Diederich: Erythropoietin, erythropoiesis and beyond. *Biochem Pharmacol* 82, 1291-1303 (2011)
DOI: 10.1016/j.bcp.2011.06.045
72. K. Maiese, Z.Z. Chong, Y.C. Shang, S. Wang: Erythropoietin: New directions for the nervous system. *Int J Mol Sci* 13, 11102-11129 (2012)
DOI: 10.3390/ijms130911102
73. A. Merelli, L. Czornyj, A. Lazarowski: Erythropoietin: A neuroprotective agent in cerebral hypoxia, neurodegeneration, and epilepsy. *Curr Pharm Des* 19, 6791-6801 (2013)
DOI: 10.2174/1381612811319380011
74. M.A. Mikati, J.A.E. Hokayem, M.E.E. Sabban: Effects of a single dose of erythropoietin on subsequent seizure susceptibility in rats exposed to acute hypoxia at P10. *Epilepsia* 48, 175-181 (2007)
DOI: 10.1111/j.1528-1167.2006.00900.x
75. R.D. Sanders, H.J. Manning, N.J. Robertson, D. Ma, A.D. Edwards, H. Hagberg, M. Maze: Preconditioning and postinsult therapies for perinatal hypoxic-ischemic injury at term. *Anesthesiology* 113, 233-249 (2010)
DOI: 10.1097/ALN.0b013e3181dc1b84
76. M.L. Brines, P. Ghezzi, S. Keenan, D. Agnello, N.C. de-Lanerolle, C. Cerami, L.M. Itri, A. Cerami: Erythropoietin crosses the blood – brain barrier to protect against experimental brain injury. *Proc Natl Acad Sci USA* 97, 10526-10531 (2000)
DOI: 10.1073/pnas.97.19.10526
77. M.G. Lykissas, A.V. Korompilias, M.D. Vekris, G.I. Mitsionis, E. Sakellariou, A.E. Beris: The role of erythropoietin in central and peripheral nerve injury. *Clin Neurol Neurosurg* 109, 639-644 (2007)
DOI: 10.1016/j.clineuro.2007.05.013
78. S.C. Sözmen, S.H. Kurul, U. Yis, K. Tugyan, B. Baykara, O. Yilmaz: Neuroprotective effects of recombinant human erythropoietin in the developing brain of rat after lithium-pilocarpine induced status epilepticus. *Brain Dev* 34, 189-195 (2012)
DOI: 10.1016/j.braindev.2011.05.002
79. D. Agnello, P. Bigini, P. Villa, T. Mennini, A. Cerami, M.L. Brines, P. Ghezzi: Erythropoietin exerts an anti-inflammatory effect on the CNS in a model of experimental autoimmune encephalomyelitis. *Brain Res* 952, 128-134 (2002)
DOI: 10.1016/S0006-8993(02)03239-0
80. M. Brines, A. Cerami: Emerging biological roles for erythropoietin in the nervous system. *Nat Rev Neurosci* 6, 484-494 (2005)
DOI: 10.1038/nrn1687

81. A. Mammis, T.K. McIntosh, A.H. Maniker: Erythropoietin as a neuroprotective agent in traumatic brain injury. *Surg Neurol* 71, 527-531 (2009)
DOI: 10.1016/j.surneu.2008.02.040
82. S. Pankratova, D. Kiryushko, K. Sonn, V. Soroka, L.B. Kohler, M. Rathje, B. Gu, K. Gotfryd, O. Clausen, A. Zharkovsky, E. Bock, V. Berezin: Neuroprotective properties of a novel, non-haematopoietic agonist of the erythropoietin receptor. *Brain* 133, 2281-2294 (2010)
DOI: 10.1093/brain/awq101
83. K.H. Jung, K. Chu, S.T. Lee, K.I. Park, J.H. Kim, K.M. Kang, S. Kim, D. Jeon, M. Kim, S.K. Lee, *et al.*: Molecular alterations underlying epileptogenesis after prolonged febrile seizure and modulation by erythropoietin. *Epilepsia* 52, 541-550 (2011)
DOI: 10.1111/j.1528-1167.2010.02916.x
84. J. Yang, Y. Huang, X. Yu, H. Sun, Y. Li, Y. Deng: Erythropoietin preconditioning suppresses neuronal death following status epilepticus in rats. *Acta Neurobiol Exp (Warsz)* 67, 141-148 (2007)
doi not found
85. U. Yis, S.H. Kurul, A. Kumral, K. Tugyan, S. Cilaker, O. Yilmaz, S. Genç, K. Genç: Effect of erythropoietin on oxygen-induced brain injury in the newborn rat. *Neurosci Lett* 448, 245-249 (2008)
DOI: 10.1016/j.neulet.2008.10.060
86. N. Seeger, C. Zellinger, A. Rode, F. Roloff, G. Bicker, V. Rusmann, S. Fischborn, H. Wendt, H. Potschka: The erythropoietin-derived peptide mimetic pHBSP affects cellular and cognitive consequences in a rat post-status epilepticus model. *Epilepsia* 52(12), 2333-2343 (2011)
DOI: 10.1111/j.1528-1167.2011.03302.x
87. K. Kobow, S. Auvin, F. Jensen, W. Löscher, I. Mody, H. Potschka, D. Prince, A. Sierra, M. Simonato, A. Pitkänen, A. Nehlig, J.M. Rho: Finding a better drug for epilepsy: Antiepileptogenesis targets. *Epilepsia* 53, 1868-1876 (2012)
DOI: 10.1111/j.1528-1167.2012.03716.x
88. E. Spandou, Z. Papadopoulou, V. Soubasi, G. Karkavelas, C. Simeonidou, A. Pazaiti, O. Guiba-Tziampiri: Erythropoietin prevents long-term sensorimotor deficits and brain injury following neonatal hypoxia-ischemia in rats. *Brain Res* 1045, 22-30 (2005)
DOI: 10.1016/j.brainres.2005.03.013
89. X.N. Wen, Y.G. Huang, J.C. Wang: Erythropoietin preconditioning on hippocampus neuronal apoptosis following status epilepticus induced by Li-pilocarpine in rats through anti-caspase-3 expression. *Neurol India* 54, 58-63 (2006)
DOI: 10.4103/0028-3886.24708
90. D. Banerjee, M. Rodriguez, M. Nag, J.W. Adamson: Exposure of endothelial cells to recombinant human erythropoietin induces nitric oxide synthase activity. *Kidney Int* 57, 1895-1904 (2000)
DOI: 10.1046/j.1523-1755.2000.00039.x
91. E. Perucca: An introduction to antiepileptic drugs. *Epilepsia* 46, 31-37 (2005)
DOI: 10.1111/j.1528-1167.2005.463007.x
92. P. Kwan, M. J. Brodie: Early identification of refractory epilepsy. *N. Engl. J. Med* 342, 314-319 (2000)
DOI: 10.1056/NEJM200002033420503
93. M.M. Acharya, B. Hattiangady, A.K. Shetty: Progress in neuroprotective strategies for preventing epilepsy. *Prog Neurobiol* 84(4), 363-404 (2008)
DOI: 10.1016/j.pneurobio.2007.10.010
94. Pitkänen: Efficacy of current antiepileptics to prevent neurodegeneration in epilepsy models. *Epilepsy Res* 50, 141-160 (2002)
DOI: 10.1016/S0920-1211(02)00076-1
95. A. O. S. Cunha, M. R. Mortari, J. L. Liberato, W. F. Santos: Neuroprotective effects of diazepam, carbamazepine, phenytoin and ketamine after pilocarpine-induced Status Epilepticus. *Basic Clin Pharmacol Toxicol* 104, 470-477 (2009)
DOI: 10.1111/j.1742-7843.2009.00403.x
96. F. Qashu, T. H. Figueiredo, V. Aroniadou-Anderjaska, J. P. Apland, M. F. M.

- Braga: Diazepam administration after prolonged status epilepticus reduces neurodegeneration in the amygdala but not in the hippocampus during epileptogenesis. *Amino Acids* 38, 189-197 (2010)
DOI: 10.1007/s00726-008-0227-2
97. A. Pitkänen, I. Kharatishvili, S. Narkilahti, K. Lukasiuk, J. Nissinen: Administration of diazepam during status epilepticus reduces development and severity of epilepsy in rat. *Epilepsy Res* 63, 27-42 (2005)
DOI: 10.1016/j.eplepsyres.2004.10.003
98. C. Brandt, A. M. Gastens, M. zhen Sun, M. Hausknecht, W. Löscher: Treatment with valproate after status epilepticus: Effect on neuronal damage, epileptogenesis, and behavioral alterations in rats. *Neuropharmacology* 51, 789-804 (2006)
DOI: 10.1016/j.neuropharm.2006.05.021
99. H. M. Capella, T. Lemos: Effect on epileptogenesis of carbamazepine treatment during the silent period of the pilocarpine model of epilepsy. *Epilepsia* 43(5), 110-111 (2002)
DOI: 10.1046/j.1528-1157.43.s.5.9.x
100. C. Brandt, M. Nozadze, N. Heuchert, M. Rattka, W. Löscher: Disease-modifying effects of phenobarbital and the NKCC1 inhibitor bumetanide in the pilocarpine model of temporal lobe epilepsy. *J Neurosci* 30, 8602-8612 (2010)
DOI: 10.1523/JNEUROSCI.0633-10.2010
101. J. Nissinen, C. H. Large, S. C. Stratton, A. Pitkänen: Effect of lamotrigine treatment on epileptogenesis: an experimental study in rat. *Epilepsy Res* 58, 119-132 (2004)
DOI: 10.1016/j.eplepsyres.2004.01.001
102. V. André, M.A. Rigoulot, E. Koning, A. Ferrandon, A. Nehlig: Long-term pregabalin treatment protects basal cortices and delays the occurrence of spontaneous seizures in the lithium-pilocarpine model in the rat. *Epilepsia* 44, 893-903 (2003)
DOI: 10.1046/j.1528-1157.2003.61802.x
103. J.B. Faure, G. Akimana, J. E. M. Carneiro, B. Cosquer, A. Ferrandon, K. Geiger, E. Koning, L. Penazzi, J.C. Cassel, A. Nehlig: A comprehensive behavioral evaluation in the lithium-pilocarpine model in rats: effects of carisbamate administration during status epilepticus. *Epilepsia* 54, 1203-1213 (2013)
DOI: 10.1111/epi.12219
104. J. François, K. Germe, A. Ferrandon, E. Koning, A. Nehlig: Carisbamate has powerful disease-modifying effects in the lithium-pilocarpine model of temporal lobe epilepsy. *Neuropharmacology* 61, 313-328 (2011)
DOI: 10.1016/j.neuropharm.2011.04.018
105. S. Chien, M. Bialer, B. Solanki, T. Verhaeghe, D. R. Dose, G. Novak, C. Yao: Pharmacokinetic interaction study between the new antiepileptic and CNS drug RWJ-333369 and carbamazepine in healthy adults. *Epilepsia* 47, 1830-1840 (2006)
DOI: 10.1111/j.1528-1167.2006.00815.x
106. Y. Liu, G. J. Yohrling, Y. Wang, T. L. Hutchinson, D. E. Brenneman, C. M. Flores, B. Zhao: Carisbamate, a novel neuromodulator, inhibits voltage-gated sodium channels and action potential firing of rat hippocampal neurons. *Epilepsy Res* 83, 66-72 (2009)
DOI: 10.1016/j.eplepsyres.2008.09.006
107. C.Y. Lee, M.L. Lee, C.C. Shih, H.H. Liou: Carisbamate (RWJ-333369) inhibits glutamate transmission in the granule cell of the dentate gyrus. *Neuropharmacology* 61, 1239-1247 (2011)
DOI: 10.1016/j.neuropharm.2011.07.022
108. S. Shim, M. El Mansari, P. Blier: Modulation of the antidepressant-like effects of sustained administration of carisbamate and lamotrigine on monoaminergic systems: electrophysiological studies in the rat brain. *J Pharmacol Exp Ther* 347, 487-496 (2013)
DOI: 10.1124/jpet.113.203315
109. H. L. Grabenstatter, F. E. Dudek: A new potential AED, carisbamate, substantially

- reduces spontaneous motor seizures in rats with kainic acid-induced epilepsy. *Epilepsia* 49, 1787-1794 (2008)
DOI: 10.1111/j.1528-1167.2008.01657.x
110. V. André, C. Dubé, J. François, C. Leroy, M.A. Rigoulot, C. Roch, I. J. Namer, A. Nehlig: Pathogenesis and pharmacology of epilepsy in the lithium-pilocarpine model. *Epilepsia* 48(15), 41-47 (2007)
DOI: 10.1111/j.1528-1167.2007.01288.x
 111. J.B. Faure, J.E. Marques-Carneiro, G. Akimana, B. Cosquer, A. Ferrandon, K. Herbeaux, E. Koning, A. Barbelivien, A. Nehlig, J.C. Cassel: Assessment of attention and executive functions in a rat model of chronic epilepsy. *Epilepsia* (In Press)
doi not found
 112. J. François, A. Boehrer, A. Nehlig: Effects of carisbamate (RWJ-333369) in two models of genetically determined generalized epilepsy, the GAERS and the audiogenic WistarAS. *Epilepsia* 49, 393-399 (2008)
DOI: 10.1111/j.1528-1167.2007.01278.x
 113. J. J. Luszczki: Third-generation antiepileptic drugs: mechanisms of action, pharmacokinetics and interactions. *Pharmacol Rep PR* 61, 197-216 (2009)
DOI: 10.1016/S1734-1140(09)70024-6
 114. M. Niebauer, M. Gruenthal: Topiramate reduces neuronal injury after experimental status epilepticus. *Brain Res* 837, 263-269 (1999)
DOI: 10.1016/S0006-8993(99)01615-7
 115. A.M. Mazarati, R.A. Baldwin, R.D. Sofia, C.G. Wasterain: Felbamate in experimental model of status epilepticus. *Epilepsia* 41, 123-127 (2000)
DOI: 10.1111/j.1528-1157.2000.tb00130.x
 116. T. Halonen, J. Nissinen, A. Pitkänen: Effect of lamotrigine treatment on status epilepticus-induced neuronal damage and memory impairment in rat. *Epilepsy Res* 46, 205-223 (2001)
DOI: 10.1016/S0920-1211(01)00278-9
 117. M. A. Rigoulot, E. Koning, A. Ferrandon, A. Nehlig: Neuroprotective properties of topiramate in the lithium-pilocarpine model of epilepsy. *J Pharmacol Exp Ther* 308, 787-795 (2004)
DOI: 10.1124/jpet.103.057091
 118. V. André, A. Ferrandon, C. Marescaux, A. Nehlig: Vigabatrin protects against hippocampal damage but is not antiepileptogenic in the lithium-pilocarpine model of temporal lobe epilepsy. *Epilepsy Res* 47, 99-117 (2001)
DOI: 10.1016/S0920-1211(01)00299-6
 119. T. Halonen, J. Nissinen, J. A. Jansen, A. Pitkänen: Tiagabine prevents seizures, neuronal damage and memory impairment in experimental status epilepticus. *Eur J Pharmacol* 299, 69-81 (1996)
DOI: 10.1016/0014-2999(95)00835-7
 120. T. Licko, N. Seeger, C. Zellinger, V. Russmann, A. Matagne, H. Potschka: Lacosamide treatment following status epilepticus attenuates neuronal cell loss and alterations in hippocampal neurogenesis in a rat electrical status epilepticus model. *Epilepsia* 54, 1176-1185 (2013)
DOI: 10.1111/epi.12196

Abbreviations: 5-HT: 5-hydroxytryptamine; 5-HT_{1A}: 5-hydroxytryptamine 1A receptor; ADK: adenosine kinase; AED: antiepileptic drug; Akt: protein kinase B; AMPA: α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor; AMY: amygdala; BBB: blood-brain barrier; CA1: *Cornus ammonis* 1; CA3: *Cornus ammonis* 3; AMPc: 3'-5'-cyclic adenosine monophosphate; CBZ: carbamazepine; CNS: central nervous system; CRS: carisbamate; DMPX: 3,7-dimethyl-1-propylxanthine; DZP: diazepam; EEG: electroencephalography; EPO: erythropoietin; EPOR: erythropoietin receptor; ERK 1: extracellular-signal-regulated kinase 1; ERK 2: extracellular-signal-regulated kinase 2; FBM: felbamate; GABA: gamma-aminobutyric acid; GABA_A: gamma-aminobutyric acid A receptor; GAERS: genetic absence epilepsy rat from Strasbourg; HIF-1: hypoxia-inducible factor-1; HPC: hippocampus; HS: hippocampal

sclerosis; IL-1 β : interleukin-1 β ; LSM: lacosamide; LTG: lamotrigine; NMDA: N-methyl-D-aspartate receptor; PGB: pregabalin; PHB: phenobarbital; pHBSP: pyroglutamate helix B surface peptides; PHT: phenytoin; PI3K: phosphatidylinositol-4,5-bisphosphate 3-kinase; PPS: perforant-path stimulation; rhEPO: recombinant human erythropoietin; R-PIA: R-N6-phenylisopropyladenosine; SCH58261: (7-(2-phenylethyl)-5-amino-2-(2-furyl)-pyrazolo(4,3-e)-1,2,4-triazolo(1,5-c) pyrimidine); SE: *status epilepticus*; SRS: spontaneous recurrent seizures; SSSE: self-sustained *status epilepticus*; SWD: spike-and-wave-discharge; TGB: tiagabine; TLE: temporal lobe epilepsy; TNF- α : tumor necrosis factor-alpha; TPM: topiramate; VGB: vigabatrin; VPA: valproate

Key Words: Temporal Lobe Epilepsy, Neuroprotection, Erythropoietin, Adenosine, Carisbamate, Review

Send correspondence to: Maria Jose da Silva Fernandes, Department of Neurology and Neurosurgery, Universidade Federal de Sao Paulo-UNIFESP, Rua Pedro de Toledo, N°669 - 2° andar, CEP 04039-032, Sao Paulo, Brazil, Tel: 5511-5576-4846, E-mail: fernandeseppm@gmail.com