

Phosphatidylinositol 3-kinase: a key regulator in adherens junction formation and function

Nathalie Rivard

Department of Anatomy and Cell Biology, Faculty of Medicine and Health Sciences, University of Sherbrooke, 3001, 12th north avenue, Sherbrooke, (Quebec), Canada, J1H 5N4

TABLE OF CONTENTS

1. Abstract
2. Introduction
3. Adherens junctions
4. Phosphatidylinositol 3-kinases
5. Regulation of E-cadherin-mediated cell-cell contacts by PI3K
6. Recruitment and activation of PI3K signaling by E-cadherin-mediated cell-cell contacts
7. PI3K is involved in E-cadherin-dependent regulation of epithelial cell differentiation and polarity
8. Conclusion
9. Acknowledgements
10. References

1. ABSTRACT

The activity of E-cadherin-adhesion complexes is under stringent control of signaling pathways. Conversely, these adhesion complexes are preferential sites for signal transduction. One class of signaling molecules reported to regulate adherens junction and to be activated by adherens junction assembly are phosphatidylinositol 3-kinases. While the exact molecular mechanisms involved are not clear, present data indicate that one of the earliest events likely involves c-Src which is rapidly activated by E-cadherin-mediated cellular aggregation and may facilitate the recruitment and activation of PI3K to E-cadherin-containing complexes. Beta-catenin, gamma-catenin and hDlg which are present at cell-cell adhesions can act as docking proteins for PI3K. Hence, cell-cell interaction leads to PtdIns(3,4,5) P_3 production in nascent cadherin contacts triggering the recruitment of proteins containing pleckstrin homology domains including the kinase PKB/Akt and the exchange factors for Rac, Tiam and Vav. PKB/Akt may be involved in the regulation of survival and proliferation while Tiam and Vav may activate Rac, resulting in reorganization of actin cytoskeleton which ultimately serves to mediate adhesive cell-cell recognition as well as epithelial cell differentiation and polarity.

2. INTRODUCTION

In epithelial tissues, the cells are tightly bound together into sheets called epithelia. These epithelia form a boundary between discrete compartments within an organism whilst also constituting the interface between the organism and the environment. The development and maintenance of epithelia require selective cell-cell and cell-extracellular matrix adhesion (reviewed in 1) which must be strong enough to maintain tissue architecture yet be sufficiently dynamic to allow essential processes such as morphogenesis or cell division to occur (reviewed in 2).

Cell adhesion is achieved by a superfamily of transmembrane glycoproteins collectively known as cell adhesion molecules and found in a diverse range of cellular junctions (reviewed in 2). A subfamily known as cadherins mediate homotypic Ca^{2+} -dependent intercellular adhesion in the majority of cell types (reviewed in 3-5) and are structurally characterized by conserved 110-amino-acid repeat sequences (cadherin repeats) present within the extracellular domain (Figure 1) (3, 4). Cadherins can be divided into four subclasses: classical cadherins, responsible for cell-cell adhesion at the zonula adherens; desmosomal cadherins, responsible for cell-cell adhesion at




Figure 1. The classic cadherin-catenin complex. The cadherin is a parallel or cis homodimer. The extracellular region of E-cadherin consists of five cadherin-type repeats (EC domains; extracellular cadherin domains) that are bound together by Ca²⁺ ions. The core-universal catenin complex consists of p120 catenin and beta-catenin which in turn binds alpha-catenin. In a less understood way, alpha-catenin binds to actin and actin-binding proteins, such as vinculin, alpha-actinin and formin-1 (32).

desmosomal junctions; and protocadherins and cadherin-related proteins whose functions are not fully elucidated (3). Individual cadherins are differentially expressed during embryonic development and mediate intercellular interactions in distinct cell types. E-cadherin is the prototypic member of the subfamily of classical cadherins and is expressed in most epithelial tissues (6). Without diminishing the importance of other cellular junctions, E-cadherin-mediated adherens junctions are particularly important in controlling the specificity, formation and maintenance of intercellular adhesion in epithelial tissues (7-11).

3. ADHERENS JUNCTIONS

Conceptually, the life of a cadherin-mediated adherens junction can be separated into three stages: formation, maintenance and disassembly (12). Adherens junctions are assembled at cell-cell contacts. During the early stages of intercellular adhesion, cells extend filopodial/lamellipodial extensions which enhance cell-cell contacts (13, 14). Such membrane protrusions involve the actin cytoskeleton and Rho GTPases including Rho, Rac and Cdc42 (15-17). Specifically, Rac has been shown to be required for efficient recruitment of F-actin to the adherens junctions in epithelial cells (18-20). Cell surface proteins such as nectins and JAMs may be involved in initial

contacts by facilitating early transient cadherin interaction and bringing membranes into apposition and/or forming signaling complexes (21-23). Cadherin-associated proteins are subsequently recruited to these nascent contacts. E-cadherin binds directly to several cytoplasmic proteins including beta-catenin and p120 (24). p120 may indirectly regulate the cadherin-actin cytoskeleton nexus by locally controlling the activity of the Rho inhibitor p190RhoGAP, hence the activation of Rho and Rac (25), as well as the rate of cadherin endocytosis (26). Beta-catenin plays a more direct role by binding to alpha-catenin (27), an actin filament binding/bundling protein that also binds other actin-binding proteins (Figure 1) (28). Clustering of adhesion molecules and recruitment of these intracellular components connecting the junctional complex to the actin cytoskeleton induce further maturation of the junction and subsequent formation of tight junctions (12, 29). However, the current view that actin filaments bind directly to alpha-catenin and therefore to E-cadherin-catenin complex was recently re-questioned (30-31). Indeed, it has been shown that alpha-catenin assembled into the cadherin-catenin complex does not bind to actin (30-31). However, when alpha-catenin dimerizes, alpha-catenin homodimers are released from the cadherin-catenin complexes and bind to actin and antagonize Arp2/3 function, inhibiting actin branching and facilitating formation of the belt of unbranched actin filaments. Alpha-catenin then acts as a molecular switch to regulate actin dynamics at adherens junctions. Alternative models could also explain the association between actin and adherens junctions. One possibility is that other molecules mediate a direct connection. One attractive candidate is the nectin-afadin system; another possibility is that the link is mediated by many weak and transient interactions (which cannot be detected biochemically) between actin binding proteins (such as zonula occludens-1, spectrin, vinculin, afadin and alpha-actinin) and adherens junction components (reviewed in 32).

The activity of E-cadherin-adhesion complexes is under stringent control of signaling pathways, many of which are dysregulated during neoplastic progression. Conversely, it is now well accepted that these adhesion complexes are also preferential sites for signal transduction. Hence, not only may signaling events regulate E-cadherin adhesive function, but cadherins themselves also participate in transducing extracellular signals to the interior. One class of signaling molecules reported to regulate adherens junction formation and to be activated by adherens junction assembly are phosphatidylinositol 3-kinases.

4. PHOSPHATIDYLINOSITOL 3-KINASES

There are three classes of phosphatidylinositol 3-kinases (PI3Ks) in mammals based on primary structure, substrate specificity and mode of regulation. Class I PI3Ks include four distinct p110 catalytic isoforms, further divided into Class IA (-alpha, -beta, -delta) and IB (-gamma); among these, p110alpha and p110beta are the isoforms mostly expressed in epithelia (reviewed in 33). *In vitro*, they are capable to convert phosphatidylinositides (PtdIns) to PtdIns-3-*P*, PtdIns-4-*P* to PtdIns(3,4)*P*₂, and




Figure 2. Biochemical activities of PI3K. The substrates (in blue) and catalytic products of class I, II and III (in purple) are indicated. The most important phosphorylation (indicated in dark purple) is of PtdIns(4,5) P_2 to PtdIns(3,4,5) P_3 which can serve as a docking site for signaling proteins with PH domains. Other inositol phospholipid kinases (and phosphatases) catalyze the reactions indicated by the black arrows.

PtdIns(4,5) P_2 to PtdIns(3,4,5) P_3 , but the *in vivo* substrate is PtdIns(4,5) P_2 (Figure 2). Class I PI3Ks primarily generate PtdIns(3,4,5) P_3 , which binds to pleckstrin homology (PH) domains and activates a variety of proteins including serine/threonine kinase PKB/Akt and exchange factors for small GTPases such as Tiam and Vav. Class IA PI3Ks are comprised of an 85kDa regulatory subunit (p85) and a 110kDa catalytic subunit (p110) (34, 35) which is activated in a phosphotyrosine-dependent manner via Src homology domain 2 (SH2 domain)-mediated recruitment to tyrosine kinase receptors or docking proteins. SH2 domains mapped within the p85 subunit specifically recognize the phosphotyrosine binding motif pYXXM (36). Class II PI3Ks have the potential to generate PtdIns(3,4,5) P_3 , although the type of PtdInsPs that they produce *in vivo* has not been fully characterized. *In vitro*, they are capable to phosphorylate PtdIns and PtdIns-4- P (Figure 2). The sole class III PI3K, Vps34, produces PtdIns-3- P (Figure 2). Class I PI3Ks as well as 3-phosphorylated PtdIns have been implicated in a number of signaling pathways in response to extracellular stimuli which in turn regulate diverse cellular programs such as cell survival, proliferation/migration, phagocytosis and glucose homeostasis (37). Class I PI3Ks have also been implicated in cancer: for example PTEN phosphatase removes the 3-phosphate from 3-phosphorylated PtdIns generated by PI3Ks and has been identified as a tumor suppressor gene mutated in a variety of human cancers (38). In addition, mutation, amplification and/or overexpression of some

PI3K Class I catalytic subunits have been observed in some carcinomas (37, 39). Furthermore, Class I PI3Ks are activated by a plethora of stimuli (36), one of which is adherens junction formation itself. Although the Class I PI3K family has been implicated in multiple cellular processes (40-42), much less is known as to the function of Class II and Class III PI3K enzymes (41, 42). In the ensuing text, PI3K will refer specifically to Class IA PI3K.

5. REGULATION OF E-CADHERIN-MEDIATED CELL-CELL CONTACTS BY PI3K

The first evidence suggesting the involvement of PI3K in adherens junction regulation in mammalian cells was reported by Hordijk and colleagues (43) who demonstrated that Tiam1 is localized at sites of cell-cell contacts in MDCK renal epithelial cells. Tiam1 is an exchange factor for Rac that requires membrane localization via its NH₂-terminal PH domain interacting with PtdIns(3,4,5) P_3 in order to activate endogenous Rac (44, 45). Overexpression of Tiam1 inhibits hepatocyte growth factor-induced cell scattering by increasing E-cadherin-mediated cell-cell adhesion. Furthermore, Tiam1/Rac signaling inhibits motile and invasive behavior of Ras-transformed MDCK cells due to restoration of E-cadherin-mediated adhesion (43). Sander and colleagues (46) subsequently confirmed that Tiam1-mediated Rac1 activation and E-cadherin-mediated cell-cell adhesion are both dependent on PI3K activity, by using pharmacological

Phosphatidylinositol 3-kinase

inhibitors of PI3Ks, namely LY294002 and wortmannin, and the dominant-negative mutant Δ p85 which carries a deletion in the binding site for the p110 catalytic subunit. Other studies further demonstrated that Rac1 is recruited at intercellular junctions during establishment of cadherin-mediated adhesion and is activated by E-cadherin-mediated homophilic interactions through PI3K (20, 47). Specifically, upon formation of nascent contacts, Rac1 becomes activated and is recruited to these sites in a PI3K-dependent manner (44-49). This in turn leads to the recruitment and activation of Arp2/3 at cadherin-mediated contacts, promoting the formation of actin branched networks (47). On the other hand, Rac1 contributes to the stable formation of nascent contacts through its effector IQGAP. By inhibiting IQGAP, Rac1 facilitates the interaction of alpha-catenin with beta-catenin (17). In *Xenopus*, IQGAP localizes at adherens junctions in regions undergoing active morphogenetic movements (50). Thus, Rac1 may regulate the balance between a dynamic to a more static state. In intestinal epithelial cells, inhibition of PI3K was also found to alter adherens junction integrity by reducing the amount of cytoskeletal-associated E-cadherin and beta-catenin at the site of cell-cell contact (51). Therefore, there appears to be a signaling hierarchy leading from PI3K to Tiam1 to Rac to the actin cytoskeleton resulting in adherens junction formation. However, the role of Tiam1 in the activation of Rac1 by E-cadherin-mediated cell-cell adhesion was recently re-questioned (52, 53). Vav2, another Rac-GDP/GTP exchange factor that localizes to the membrane via its NH₂-terminal PH domain interacting with $PtdIns(3,4,5)P_3$, has recently been shown to be necessary for PI3K-dependent activation of Rac1 during adherens junction formation (53). Interestingly, Vav2 is also recruited to E-cadherin-based cell-cell adhesion sites where it is tyrosine phosphorylated and activated by c-Src (53).

Recent evidences suggest that the recruitment and activation of PI3K and Rac1 upon E-cadherin engagement are, however, transient. Indeed, it has been reported that Rac1 tends to localize at the dynamic, expanding edges of a forming contact, but not when contacts had already formed and the membranes had become more quiescent (54). Moreover, direct analysis of subcellular sites of Rac1 activation between contacting MDCK using the Raichu FRET-based biosensors (55, 56) revealed that Rac1 activity is high in a restricted zone at the edges of expanding cell-cell contacts but is very low within the established cell-cell contact where E-cadherin has accumulated. More recently, a study from Nelson's laboratory describes the spatiotemporal immediate-early signaling induced by E-cadherin engagement and adhesion (57). By using laser tweezers to place beads coated with functional E-cadherin extracellular domain on cells, they quantified the kinetics of E-cadherin, PI3K and Rac1 distributions immediately following E-cadherin engagement. They found that cellular E-cadherin accumulates rapidly around beads and phosphoinositides and Rac1 co-accumulate with E-cadherin, reach peak levels with E-cadherin but then rapidly disperse. E-cadherin accumulation was dependent on membrane dynamics and actin polymerization, but actin did not stably co-accumulate

with E-cadherin. Thus, the recruitment of PI3K and Rac1 upon E-cadherin engagement appears to be transient. They propose a model in which initial E-cadherin accumulation requires active membrane dynamics and involves diffusion-mediated trapping at contact sites; to propagate further contacts, PI3K and Rac1 are transiently activated by E-cadherin engagement and initiate a new round of membrane dynamics, but they are subsequently suppressed at that site to allow maintenance and strengthening of weak E-cadherin-mediated adhesion (57).

6. RECRUITMENT AND ACTIVATION OF PI3K SIGNALING BY E-CADHERIN-MEDIATED CELL-CELL CONTACTS

Pece and colleagues (58) were the first to assess whether adherens junction assembly can affect the state of activation of PI3K signaling. As epithelial cells require Ca^{2+} in order to form homophilic cell-cell adhesions, a simple method to study the adhesive properties of surface molecules involves the disruption of intercellular Ca^{2+} -dependent homotypic boundaries by EGTA-treatment and the re-establishment of cell-cell contacts by the subsequent restoration of Ca^{2+} ions (so-called Ca^{2+} -switch) (59). By using Ca^{2+} -switch experiments, Pece and colleagues demonstrated that engagement of E-cadherins in homophilic adhesion with neighboring cells promotes a substantial activation of PI3K-dependent PKB/Akt and the rapid translocation of PKB/Akt to the nucleus (58). Such stimulation of PI3K by E-cadherin-mediated cell-cell contact has also been reported in other epithelial cell types (51, 60). Furthermore, a physical association between PI3K and E-cadherin-containing multiprotein complexes is observed in response to cell-cell contact formation, thus providing a likely mechanism for PKB/Akt activation (51, 61, 62). However, the interaction between E-cadherin and p85/PI3K is indirect and may involve intermediate proteins (61, 62).

The molecular mechanisms by which PI3K is recruited and activated to E-cadherin-mediated cell-cell contacts have been studied to some extent, although the exact sequence of molecular events involved in this process is still not clear. As in many other models of PI3K signaling, recruitment of PI3K to the plasma membrane is a key step in the cadherin-activated signaling pathway. In many instances of receptor-activated PI3K signaling, binding of p85 is in itself a response to upstream tyrosine kinase (reviewed in 63). Recruitment commonly entails high affinity interactions between SH2 domains and tyrosine-phosphorylated sequences in components of the receptor complex (64, 65). Many tyrosine kinases are known to concentrate at cadherin-adhesive contacts (66-68) while c-src has been shown to co-recruit to cadherin homophilic contacts and to be critical for the recruitment and activation of PI3K to E-cadherin (69, 70).

Three cadherin-associated proteins have been reported to bind the p85 subunit of PI3K at E-cadherin-mediated cell-cell contacts (Figure 3), namely beta-catenin (61, 71, 72), gamma-catenin (73) and hDlg (62). Beta-catenin has been shown to directly interact with the N-

Phosphatidylinositol 3-kinase

terminal SH2 domain of p85 in the cadherin-based adhesion complex of human keratinocytes and MDCK epithelial cells. However, this interaction does not require tyrosine phosphorylation (61) and thus the mechanism by which beta-catenin interacts with p85 needs to be clarified.

Gamma-catenin may also represent an additional docking protein for PI3K within E-cadherin complexes. This protein is known to co-precipitate with the p110alpha catalytic subunit in keratinocytes and demonstrates prominent reactivity to an antibody recognizing phosphorylated YXXM motifs (73). In parallel, tyrosine kinase inhibition, in addition to suppressing the activation of PI3K pathway, also blocks the association of PI3K with E-cadherin-mediated complexes and the reactivity of gamma-catenin to the PI3K docking site antibody (73).

hDlg, a member of the membrane-associated guanylate kinase (MAGUK)-like protein family, has been shown to be involved in PI3K recruitment and binding to E-cadherin adhesion complex (62). Interestingly, members of the MAGUK-like protein family contribute to proper junction and cell polarity (74-76). DLG-1, a *C. elegans* homologue of the *Drosophila* disc-large (Dlg), is required for the proper assembly and function of adherens junctions (77-79). Indeed, Firestein and Rongo (79) have shown that DLG-1 is predominantly localized at the adherens junctions of the epidermis, intestine and pharynx of embryonic and adult nematodes. Conversely, DLG-1-deficient embryos exhibit abnormal adherens junction formation and disorganization of the actin cytoskeleton (79). Human Dlg (hDlg) contains many protein-protein interaction domains including three PDZ (PSD-95/Dlg/ZO-1) domains, e.g. a SH3 domain, a GUK domain and alternatively spliced insertions (80-82). Consistent with the cellular localization of Dlg to lateral junctions (77), hDlg localizes to regions of cell-cell contact in human epithelial cells (62, 83, 84). We have recently reported that hDlg is required for the recruitment and maintenance of p85/PI3K to the E-cadherin adhesion complex (62). Indeed, epithelial cells containing low hDlg levels fail to recruit PI3K to E-cadherin-mediated cell-cell contacts and are unable to organize their cortical actin cytoskeleton. PI3K recruitment together with an organized cortical actin cytoskeleton are hence indispensable for proper adherens junction formation and stabilization as well as for morphological and functional epithelial differentiation (51). In addition to demonstrating a role for hDlg in the regulation of adherens junction assembly and function, our recent data also provide insights into how hDlg recruits PI3K signaling to the E-cadherin adhesion complex (62). Firstly, the direct interaction between p85/PI3K and hDlg is mediated by both SH2 domains of p85/PI3K. Accordingly, tyrosine phosphorylation of hDlg is necessary for p85/hDlg association. Since the binding of both SH2 domains of p85 is required for full activation of PI3K (63), this suggests that hDlg/p85 interaction may contribute to PI3K activation. Preliminary experiments demonstrate that hDlg is indeed tyrosine phosphorylated in intestinal epithelial cells (N. Rivard, unpublished data). However, the tyrosine kinase(s) which phosphorylate(s) hDlg in epithelial cells

is(are) unknown, although c-Src may constitute a plausible candidate (69, 70). Secondly, Far-Western analysis demonstrates that the regulatory p85 subunit of PI3K directly interacts with specific variants of hDlg and more specifically with hypo-phosphorylated forms (fast migrating bands) present in differentiated polarized intestinal and renal epithelial cells. Interestingly, the dephosphorylation of hDlg on serine and threonine residues by PP1 phosphatase clearly reveals an interaction of the SH2 domains of p85 with the fast migrating forms of hDlg, suggesting that in undifferentiated subconfluent non polarized cells, hDlg is already tyrosine phosphorylated but its serine/threonine phosphorylation prevents its association with p85/PI3K (62). Therefore, it appears that the p85 binding site on hDlg protein is only unmasked upon confluency and differentiation. The change in hDlg phosphorylation state on serine and threonine during differentiation, together with the association of p85 with hypo-phosphorylated forms of hDlg, are consistent with the hypothesis that serine/threonine phosphorylation negatively regulates the association of p85/PI3K with hDlg to the E-cadherin adhesion complex and hence its scaffolding and clustering activity. In this respect, several phosphorylation consensus sequences for various serine/threonine kinases are distributed throughout the hDlg protein. However, further studies are required at this point to identify the actual phosphorylated serine and/or threonine on hDlg and their role in hDlg function in epithelial cells as well as to clarify the mechanism by which hDlg phosphorylation specifically inhibits p85/PI3K association.

7. PI3K IS INVOLVED IN E-CADHERIN-DEPENDENT REGULATION OF EPITHELIAL CELL DIFFERENTIATION AND POLARITY

In epithelial cells in culture, the localization and the cellular effect of PI3K activation clearly differ depending on the conditions in which cells are maintained, and in particular on the degree of confluence. For example, in intestinal epithelial cells, the p85 regulatory subunit of PI3K exhibits a cytoplasmic distribution in subconfluent growing cells, probably associated with Focal Adhesion kinase (FAK) (85) or other proteins, while a localized distribution pattern to cell-cell interfaces has been observed in confluent differentiating cells (51). Co-immunoprecipitation experiments confirmed the significant association of E-cadherin with p85/PI3K only in confluent differentiating intestinal epithelial cells (51). Accordingly, Tiam1 is localized in lamellipodia and membrane ruffles of migrating fibroblastoid cells while it is found in the adherens junctions of non-motile epithelial monolayers (46). Hence, when localized into the cytoplasm, PI3K is involved in cell migration (86-91) and proliferation (reviewed in 37, 38) probably by its association with FAK or other partners whereas PI3K may control not only adherens junction assembly but also differentiation and polarity when it is translocated to E-cadherin-mediated cell-cell contacts at confluency. Indeed, E-cadherin-mediated cell-cell attachment plays an important role in the morphological and functional differentiation of many epithelia (92-98). One of the key questions in epithelial

development is what triggers the differentiation process. In this regard, it has been demonstrated that cell-cell junction systems, particularly adherens junctions, play a critical role in the control of cell differentiation during ontogeny as well as during the continuous renewal of epithelial cells in the mature organ. For instance, studies with E-cadherin knockout mice have revealed that E-cadherin-mediated cell adhesion is essential for the compaction of mesenchymal cells and their transition to a polarized epithelium (99, 100). In a chimeric-transgenic animal model, expression of a dominant-negative N/E-cadherin mutant in villous enterocytes resulted in perturbation of cell-cell adhesion associated with an increased enterocyte migration rate along the crypt-villus axis, the loss of the differentiated, polarized phenotype and increased apoptosis (101). The tissue-specific ablation of E-cadherin impairs the differentiation of both epidermal keratinocytes and mammary epithelial cells (102-104). However, signaling components that relay the signal from E-cadherin to nuclear targets for the control of epithelial tissue-specific gene expression remain elusive. The fact that PI3K is recruited to and activated by E-cadherin-mediated cell-cell contacts in epithelial cells suggests that it may constitute one of these signaling components. For instance, inhibition of PI3K activity has been shown to repress the expression of intestine-specific genes such as sucrase-isomaltase and to delay functional and morphological differentiation of enterocytes (51). In addition, blockade of PI3K inhibits the expression of late differentiation markers such as loricrin and filaggrin in epidermal keratinocytes (73). PI3K is also involved in 3-dimensional morphogenesis and tissue-specific differentiation in the mammary gland (104). Hence, PI3K plays a primary role in regulating the integrity of adherens junctions, which in turn appears crucial for the efficient functional and morphological differentiation of epithelial cells. Conversely, PI3K may act as an intermediate in the formation of adherens junctions and differentiation, suggesting a bi-directional regulation between PI3K activity and adherens junction assembly such as previously observed for Rac and Cdc42 (20, 105). The manner in which PI3K relays the signal from E-cadherin to nuclear targets for the control of epithelial tissue-specific gene expression remains to be further clarified however. In intestinal epithelial cells, one of the downstream molecular events resulting from E-cadherin-mediated cellular aggregation is the activation of the mitogen-activated protein kinase p38alpha (Figure 4) (51), which controls differentiation of enterocytes (106); indeed, the stimulation of intestine-specific gene expression by p38alpha is mediated by the Cdx2 transcription factor, a p38 substrate known to be essential in enterocyte differentiation (106). In addition to regulating epithelial differentiation, E-cadherin-dependent PI3K activation may have other unexpected roles in the biology of epithelial cells. Indeed, we have demonstrated that E-cadherin-mediated cell-cell adhesion triggers PKB/Akt-induced inhibition of Raf-1/MEK/ERK cascade in a PI3K-dependent manner (Figure 4) (107). Strong evidences exist for the critical involvement of the Raf-1/MEK/ERK signaling cascade in the regulation of intestinal epithelial cell proliferation (108, 109). Hence,

PI3K may play a role in the transition between proliferation and differentiation in the intestinal epithelium, a role shared by E-cadherin (10, 11).

Importantly, $\text{PtdIns}(3,4,5)P_3$ has recently emerged as a key determinant of epithelial polarity. For example, in chemotaxing cells, $\text{PtdIns}(3,4,5)P_3$ accumulates at the leading edge of migrating cells (110). This results in the recruitment of many PH domain-containing proteins which in turn produce the polarized phenotype and motility of chemotaxing cells. In polarized epithelial cells, $\text{PtdIns}(3,4,5)P_3$ is stably localized at the basolateral membrane and is excluded from the apical plasma membrane (111, 112). Recently, Mostov and colleagues reported that $\text{PtdIns}(3,4,5)P_3$ is a key signal for the formation and expansion of the basolateral surface (112). In their experiments, exogenous $\text{PtdIns}(3,4,5)P_3$ ectopically implanted into the apical surface constituted a sufficient signal to transform the composition of the apical surface to a basolateral type. Interestingly, inhibition of endogenous $\text{PtdIns}(3,4,5)P_3$ production by LY294002 causes a decrease in the amount of lateral surface, resulting in shorter cells (51, 112). Hence, E-cadherin-mediated cell-cell interaction (and perhaps integrin-mediated cell-matrix interaction), by activating PI3K, maintains $\text{PtdIns}(3,4,5)P_3$ production at the basolateral membrane (Figure 4). The mechanism by which a gradient of this freely diffusible lipid is maintained may also involve PTEN which strongly localizes to the apical plasma membrane. In fact, PTEN segregates $\text{PtdIns}(4,5)P_2$ to the apical surface, recruiting annexin 2 and Cdc42 which spatially regulate actin assembly. In turn, Cdc42 goes on to recruit the Par6/aPKC complex, which further stabilizes axial polarity (113).

8. CONCLUSION

The studies outlined above illustrate the significant progress achieved to date in understanding the importance of the PI3K signaling network in adherens junction formation and signaling function (Figure 4). However, numerous questions still remain unanswered. The upstream mechanisms that allow cadherin adhesion to modulate PI3K activity remain to be clarified. Obviously, one of the earliest events likely involves c-Src which is rapidly activated by E-cadherin-mediated cellular aggregation and may facilitate the recruitment of PI3K to E-cadherin-containing complexes. Whether this reflects a role of c-Src in regulating the binding of PI3K to the cadherin-catenin complex and/or catalytic activation of PI3K by c-Src remains to be determined. PI3K may not be the only target for cadherin-activated c-Src signaling as other potential c-Src substrates such as p120, beta-catenin, gamma-catenin, cortactin and hDlg are also present at cell-cell adhesions. As they become tyrosine-phosphorylated, some of these proteins may provide binding sites for PI3K. Whether the upstream docking protein for PI3K varies with differences in cellular background or context also needs to be addressed. It will be important in the future to understand in detail, how phosphorylation of the cadherin/catenin complex and of other substrates in its




Figure 4. Outside-in signaling, receptor-like function of E-cadherin involving PI3K. After primary adhesion (a), PI3K is recruited to E-cadherin and activated. PtdIns(3,4,5) P_3 may participate to the recruitment of PH-containing proteins including the Rac exchange factors, Tiam and Vav2 (b), and the protein kinases PKB/Akt and PDK1 (not shown). Vav2 and Tiam promote Rac1 activation which, as recently reported (54, 57), stimulates membrane (c) and actin dynamics (d) adjacent to the initial site of contact, increasing the probability of additional, new E-cadherin engagements (e) (57). Alpha-catenin dimerizes, alpha-catenin homodimers are released from the cadherin-catenin complexes, bind to actin (f) and antagonize Arp2/3 function, inhibiting actin branching and facilitating formation of the belt of unbranched actin filaments. Alternatively, other actin-binding proteins such as vinculin, afadin (not shown) and alpha-actinin may provide a link with actin cytoskeleton. Meanwhile, accumulation of PtdIns(3,4,5) P_3 in the membrane signals for the formation and expansion of the basolateral surface (g) (112) and Rac1 promotes orientation of polarity and lumen formation (h) (114). In addition, in specific cellular contexts, Rac1 triggers the activation of downstream signaling effectors such as the mitogen-activated protein kinase p38alpha (i) while PKB/Akt leads to the inhibition of ERK signaling cascade (j), hence promoting cell cycle arrest of confluent epithelial cells. PtdIns(3,4,5) P_3 -dependent activation of PKB/Akt may also promote survival of polarized epithelial cells (k).

close vicinity translates into changes in adhesive strength and downstream signaling. Finally, more studies are needed to understand how the spatial and temporal expression of PI3K activity intersects with cellular context, polarity determinants and the signaling of diverse receptors, including cell-cell adhesive receptors such as E-cadherin.

9. ACKNOWLEDGEMENTS

I thank Pierre Pothier for the critical reading of the manuscript. Some of the work reported in this manuscript was supported by a grant from Canadian Institutes of Health Research (MT-14405). Nathalie Rivard

Phosphatidylinositol 3-kinase

is a recipient of a Canadian Research Chair in Signaling and Digestive Physiopathology.

10. REFERENCES

1. X. Chen and B.M. Gumbiner: Crosstalk between different adhesion molecules. *Curr Opin Cell Biol* 18, 572-578 (2006)
2. E. Knust: Regulation of epithelial cell shape and polarity by cell-cell adhesion. *Mol Membr Biol* 19, 113-120 (2002)
3. J.M. Halbleib and W.J. Nelson: Cadherins in development: cell adhesion, sorting, and tissue morphogenesis. *Genes Dev* 20, 3199-3214 (2006)
4. S. Pokutta and W.I. Weis: Structure and mechanism of cadherins and catenins in cell-cell contacts. *Annu Rev Cell Dev Biol* 23, 237-261 (2007)
5. A.S. Yap, M.S. Crampton and J. Hardin: Making and breaking contacts: the cellular biology of cadherin regulation. *Curr Opin Cell Biol* 9, 508-514 (2007)
6. S.D. Patel, C.P. Chen, F. Bahna, B. Honig and L. Shapiro: Cadherin-mediated cell-cell adhesion: sticking together as a family. *Curr Opin Struct Biol* 13, 690-698 (2003)
7. B. Gumbiner, B. Stevenson and A. Grimaldi: The role of the cell adhesion molecule uvomorulin in the formation and maintenance of the epithelial junctional complex. *J Cell Biol* 107, 1575-1587 (1988)
8. J.E. Lewis, P.J. Jensen and M.J. Wheelock: Cadherin function is required for human keratinocytes to assemble desmosomes and stratify in response to calcium. *J Invest Dermatol* 102, 870-877 (1994)
9. M. Takeichi: Cadherins: a molecular family important in selective cell-cell adhesion. *Annu Rev Biochem* 59, 237-252 (1990)
10. M.L. Hermiston and J.I. Gordon: *In vivo* analysis of cadherin function in the mouse intestinal epithelium: essential roles in adhesion, maintenance of differentiation, and regulation of programmed cell death. *J Cell Biol* 129, 489-506 (1995)
11. M.L. Hermiston, M.H. Wong and J.I. Gordon: Forced expression of E-cadherin in the mouse intestinal epithelium slows cell migration and provides evidence for nonautonomous regulation of cell fate in a self-renewing system. *Genes Dev* 15, 985-996 (1996)
12. M. Perez-Moreno and E. Fuchs: Catenins: keeping cells from getting their signals crossed. *Dev Cell* 11, 601-612 (2006)
13. C.L. Adams, Y.T. Chen, S.J. Smith and W.J. Nelson: Mechanisms of epithelial cell-cell adhesion and cell compaction revealed by high-resolution tracking of E-cadherin-green fluorescent protein. *J Cell Biol* 142, 1105-1119 (1998)
14. V. Vasioukhin, C. Bauer, M. Yin and E. Fuchs: Directed actin polymerization is the driving force for epithelial cell-cell adhesion. *Cell* 100, 209-219 (2000)
15. S. Etienne-Manneville and A. Hall: Rho GTPases in cell biology. *Nature* 420, 629-635 (2002)
16. S. Kuroda, M. Fukata, K. Fujii, T. Nakamura, I. Izawa and K. Kaibuchi: Regulation of cell-cell adhesion of MDCK cells by Cdc42 and Rac1 small GTPases. *Biochem Biophys Res Commun* 240, 430-435 (1997)
17. S. Kuroda, M. Fukata, M. Nakagawa, K. Fujii, T. Nakamura, T. Ookubo, I. Izawa, T. Nagase, N. Nomura, H. Tani, I. Shoji, Y. Matsuura, S. Yonehara and K. Kaibuchi: Role of IQGAP1, a target of the small GTPases Cdc42 and Rac1, in regulation of E-cadherin-mediated cell-cell adhesion. *Science* 281, 832-835 (1998)
18. V.M. Braga, L.M. Machesky, A. Hall and N.A. Hotchin: The small GTPases Rho and Rac are required for the establishment of cadherin-dependent cell-cell contacts. *J Cell Biol* 137, 1421-1431 (1997)
19. K. Takaishi, T. Sasaki, H. Kotani, H. Nishioka and Y. Takai: Regulation of cell-cell adhesion by rac and rho small G proteins in MDCK cells. *J Cell Biol* 139, 1047-1059 (1997)
20. M. Nakagawa, M. Fukata, M. Yamaga, N. Itoh and K. Kaibuchi: Recruitment and activation of Rac1 by the formation of E-cadherin-mediated cell-cell adhesion sites. *J Cell Sci* 114, 1829-1838 (2001)
21. A. Fukuhara, K. Shimizu, T. Kawakatsu, T. Fukuhara and Y. Takai: Involvement of nectin-activated Cdc42 small G protein in organization of adherens and tight junctions in Madin-Darby canine kidney cells. *J Biol Chem* 278, 51885-51893 (2003)
22. Y. Takai, K. Irie, K. Shimizu, T. Sakisaka and W. Ikeda: Nectins and nectin-like molecules: roles in cell adhesion, migration, and polarization. *Cancer Sci* 94, 655-667 (2003)
23. K. Tachibana, H. Nakanishi, K. Mandai, K. Ozaki, W. Ikeda, Y. Yamamoto, A. Nagafuchi, S. Tsukita and Y. Takai: Two cell adhesion molecules, nectin and cadherin, interact through their cytoplasmic domain-associated proteins. *J Cell Biol* 150, 1161-1176 (2000)
24. W.I. Weis and W.J. Nelson: Re-solving the cadherin-catenin-actin conundrum. *J Biol Chem* 281, 35593-35597 (2006)
25. P.Z. Anastasiadis and A.B. Reynolds: Regulation of Rho GTPases by p120-catenin. *Curr Opin Cell Biol* 13, 604-610 (2001)

Phosphatidylinositol 3-kinase

26. M.A. Davis, R.C. Ireton and A.B. Reynolds: A core function for p120-catenin in cadherin turnover. *J Cell Biol* 163, 525-534 (2003)
27. H. Aberle, S. Butz, J. Stappert, H. Weissig, R. Kemler and H. Hoschuetzky: Assembly of the cadherin-catenin complex *in vitro* with recombinant proteins. *J Cell Sci* 107, 3655-3663 (1994)
28. A. Kobiela and E. Fuchs: Alpha-catenin: at the junction of intercellular adhesion and actin dynamics. *Nat Rev Mol Cell Biol* 5, 614-625 (2004)
29. M.R. Kooistra, N. Dubé and J.L. Bos: Rap1: a key regulator in cell-cell junction formation. *J Cell Sci* 120, 17-22 (2007)
30. F. Drees, S. Pokutta, S. Yamada, W.J. Nelson and W.I. Weis: Alpha-catenin is a molecular switch that binds E-cadherin-beta-catenin and regulates actin-filament assembly. *Cell* 123, 903-915 (2005)
31. S. Yamada, S. Pokutta, F. Drees, W.I. Weis and W.J. Nelson: Deconstructing the cadherin-catenin-actin complex. *Cell* 123, 889-901 (2005)
32. J. Gates and M. Peifer: Can 1000 reviews be wrong? Actin, alpha-Catenin, and adherens junctions. *Cell* 123, 769-772 (2005)
33. K. E. Anderson and S.P. Jackson: Class I phosphoinositide 3-kinases. *Int J Biochem Cell Biol* 35, 1028-1033 (2003)
34. M. Otsu, I. Hiles, I. Gout, M.J. Fry, F. Ruiz-Larrea, G. Panayotou, A. Thompson, R. Dhand, J. Hsuan, N. Totty, A.D. Smith, S.J. Morgan, S.A. Courtneidge, P.J. Parker and M.D. Waterfield: Characterization of two 85 kd proteins that associate with receptor tyrosine kinases, middle-T/pp60c-src complexes, and PI3-kinase. *Cell* 65, 91-104 (1991)
35. I.D. Hiles, M. Otsu, S. Volinia, M.J. Fry, I. Gout, R. Dhand, G. Panayotou, F. Ruiz-Larrea, A. Thompson, N.F. Totty, J.J. Hsuan, S.A. Courtneidge, P.J. Parker and M.D. Waterfield: Phosphatidylinositol 3-kinase: structure and expression of the 110 kd catalytic subunit. *Cell* 70, 419-429 (1992)
36. L.C. Cantley, K.R. Auger, C. Carpenter, B. Duckworth, A. Graziani, R. Kapeller and S. Soltoff: Oncogenes and signal transduction. *Cell* 64, 281-302 (1991)
37. R. Katso, K. Okkenhaug, K. Ahmadi, S. White, J. Timms and M.D. Waterfield: Cellular function of phosphoinositide 3-kinases: implications for development, homeostasis, and cancer. *Annu Rev Cell Dev Biol* 17, 615-675 (2001)
38. K.M. Yamada and M. Araki: Tumor suppressor PTEN: modulator of cell signaling, growth, migration and apoptosis. *J Cell Sci* 114, 2375-2382 (2001)
39. B. Vogelstein and K.W. Kinzler: Cancer genes and the pathways they control. *Nat Med* 10, 789-799 (2004)
40. K. Okkenhaug and B. Vanhaesebroeck: New responsibilities for the PI3K regulatory subunit p85 alpha. *Sci STKE* 65, PE1 (2001)
41. B. Vanhaesebroeck, S.J. Leever, K. Ahmadi, J. Timms, R. Katso, P.C. Driscoll, R. Woscholski, P.J. Parker and M.D. Waterfield: Synthesis and function of 3-phosphorylated inositol lipids. *Annu Rev Biochem* 70, 535-602 (2001)
42. B. Vanhaesebroeck, G.E. Jones, W.E. Allen, D. Zicha, R. Hooshmand-Rad, C. Sawyer, C. Wells, M.D. Waterfield and A.J. Ridley: Distinct PI(3)Ks mediate mitogenic signaling and cell migration in macrophages. *Nat Cell Biol* 1, 69-71 (1999)
43. P.L. Hordijk, J.P. ten Klooster, R.A. van der Kammen, F. Michiels, L.C. Oomen and J.G. Collard: Inhibition of invasion of epithelial cells by Tiam1-Rac signaling. *Science* 278, 1464-1466 (1997)
44. F. Michiels, J.C. Stam, P.L. Hordijk, R.A. van der Kammen, L. Ruuls-Van Stalle, C.A. Feltkamp and J.G. Collard: Regulated membrane localization of Tiam1, mediated by the NH2-terminal pleckstrin homology domain, is required for Rac-dependent membrane ruffling and C-Jun NH2-terminal kinase activation. *J Cell Biol* 137, 387-398 (1997)
45. J.C. Stam, E.E. Sander, F. Michiels, F.N. van Leeuwen, H.E. Kain, R.A. van der Kammen and J.G. Collard: Targeting of Tiam1 to the plasma membrane requires the cooperative function of the N-terminal pleckstrin homology domain and an adjacent protein interaction domain. *J Biol Chem* 272, 28447-28454 (1997)
46. E.E. Sander, S. van Delft, J.P. ten Klooster, T. Reid, R.A. van der Kammen, F. Michiels and J.G. Collard: Matrix-dependent Tiam1/Rac signaling in epithelial cells promotes either cell-cell adhesion or cell migration and is regulated by phosphatidylinositol 3-kinase. *J Cell Biol* 143, 1385-1398 (1998)
47. E.M. Kovacs, M. Goodwin, R.G. Ali, A.D. Paterson and A.S. Yap: Cadherin-directed actin assembly: E-cadherin physically associates with the Arp2/3 complex to direct actin assembly in nascent adhesive contacts. *Curr Biol* 12, 379-382 (2002)
48. J.S. Ehrlich, M.D. Hansen and W.J. Nelson: Spatio-temporal regulation of Rac1 localization and lamellipodia dynamics during epithelial cell-cell adhesion. *Dev Cell* 3, 259-270 (2002)
49. N.K. Noren, C.M. Niessen, B.M. Gumbiner and K. Burridge: Cadherin engagement regulates Rho family GTPases. *J Biol Chem* 276, 33305-33308 (2001)

Phosphatidylinositol 3-kinase

50. S. Yamashiro, T. Noguchi and I. Mabuchi: Localization of two IQGAPs in cultured cells and early embryos of *Xenopus laevis*. *Cell Motil Cytoskeleton* 55, 3 6-50 (2003)
51. P. Laprise, P. Chailier, M. Houde, J.F. Beaulieu, M.J. Boucher and N. Rivard: Phosphatidylinositol 3-kinase controls human intestinal epithelial cell differentiation by promoting adherens junction assembly and p38 MAPK activation. *J Biol Chem* 277, 8226-8234 (2002)
52. A. Kraemer, M. Goodwin, S. Verma, A.S. Yap and R.G. Ali: Rac is a dominant regulator of cadherin-directed actin assembly that is activated by adhesive ligation independently of Tiam1. *Am J Physiol Cell Physiol* 292, C1061-1069 (2007)
53. T. Fukuyama, H. Ogita, T. Kawakatsu, M. Inagaki and Y. Takai: Activation of Rac by cadherin through the c-Src-Rap1-phosphatidylinositol 3-kinase-Vav2 pathway. *Oncogene* 25, 8-19 (2006)
54. S. Yamada and W.J. Nelson: Localized zones of Rho and Rac activities drive initiation and expansion of epithelial cell-cell adhesion. *J Cell Biol* 178, 517-527 (2007)
55. R.E. Itoh, K. Kurokawa, Y. Ohba, H. Yoshizaki, N. Mochizuki and M. Matsuda: Activation of rac and cdc42 video imaged by fluorescent resonance energy transfer-based single-molecule probes in the membrane of living cells. *Mol Cell Biol* 22, 6582-6591 (2002)
56. H. Yoshizaki, Y. Ohba, K. Kurokawa, R.E. Itoh, T. Nakamura, N. Mochizuki, K. Nagashima, and M. Matsuda: Activity of Rho-family GTPases during cell division as visualized with FRET-based probes. *J Cell Biol* 162, 223-232 (2003)
57. T.D. Perez, T. Tamada, M.P. Sheetz and W.J. Nelson: Immediate-early signaling induced by E-cadherin engagement and adhesion. *J Biol Chem* (2007) [Epub ahead of print]
58. S. Pece, M. Chiariello, C. Murga and J.S. Gutkind: Activation of the protein kinase Akt/PKB by the formation of E-cadherin-mediated cell-cell junctions. Evidence for the association of phosphatidylinositol 3-kinase with the E-cadherin adhesion complex. *J Biol Chem* 274, 19347-19351 (1999)
59. T. Volberg, B. Geiger, J. Kartenbeck and W.W. Franke: Changes in membrane-microfilament interaction in intercellular adherens junctions upon removal of extracellular Ca²⁺ ions. *J Cell Biol* 102, 1832-1842 (1986)
60. E. Bergin, J.S. Levine, J.S. Koh and W. Lieberthal: Mouse proximal tubular cell-cell adhesion inhibits apoptosis by a cadherin-dependent mechanism. *Am J Physiol Renal Physiol* 278, F758-768 (2000)
61. R.J. Woodfield, M.N. Hodgkin, N. Akhtar, M.A. Morse, K.J. Fuller, K. Saqib, N.T. Thompson and M.J. Wakelam: The p85 subunit of phosphoinositide 3-kinase is associated with beta-catenin in the cadherin-based adhesion complex. *Biochem J* 360, 335-344 (2001)
62. P. Laprise, A. Viel and N. Rivard: Human homolog of disc-large is required for adherens junction assembly and differentiation of human intestinal epithelial cells. *J Biol Chem* 279, 10157-10166 (2004)
63. M.P. Wymann and L. Pirola: Structure and function of phosphoinositide 3-kinases. *Biochim Biophys Acta* 1436, 127-150 (1998)
64. S. Dowler, L. Montalvo, D. Cantrell, N. Morrice and D.R. Alessi: Phosphoinositide 3-kinase-dependent phosphorylation of the dual adaptor for phosphotyrosine and 3-phosphoinositides by the Src family of tyrosine kinase. *Biochem J* 349, 605-610 (2000)
65. B. Vanhaesebroeck, S.J. Leever, K. Ahmadi, J. Timms, R. Katso, P.C. Driscoll, R. Woscholski, P.J. Parker and M.D. Waterfield: Synthesis and function of 3-phosphorylated inositol lipids. *Annu Rev Biochem* 70, 535-602 (2001)
66. S. Tsukita, K. Oishi, T. Akiyama, Y. Yamanashi, T. Yamamoto and S. Tsukita: Specific proto-oncogenic tyrosine kinases of src family are enriched in cell-to-cell adherens junctions where the level of tyrosine phosphorylation is elevated. *J Cell Biol* 113, 867-879 (1991)
67. E. Calautti, S. Cabodi, P.L. Stein, M. Hatzfeld, N. Kedersha and G.P. Dotto: Tyrosine phosphorylation and src family kinases control keratinocyte cell-cell adhesion. *J Cell Biol* 141, 1449-1465 (1998)
68. E. Calautti, M. Grossi, C. Mammucari, Y. Aoyama, M. Pirro, Y. Ono, J. Li and G.P. Dotto: Fyn tyrosine kinase is a downstream mediator of Rho/PRK2 function in keratinocyte cell-cell adhesion. *J Cell Biol* 156, 137-148 (2002)
69. R.W. McLachlan, A. Kraemer, F.M. Helwani, E.M. Kovacs and A.S. Yap: E-cadherin adhesion activates c-Src signaling at cell-cell contacts. *Mol Biol Cell* 18, 3214-3223 (2007)
70. J.H. Pang, A. Kraemer, S.J. Stehbens, M.C. Frame and A.S. Yap: Recruitment of phosphoinositide 3-kinase defines a positive contribution of tyrosine kinase signaling to E-cadherin function. *J Biol Chem*, 280, 3043-3050 (2005)
71. P. Carmeliet, M.G. Lampugnani, L. Moons, F. Breviario, V. Compernelle, F. Bono, G. Balconi, R. Spagnuolo, B. Oostuyse, M. Dewerchin, A. Zanetti, A. Angellilo, V. Mattot, D. Nuyens, E. Lutgens, F. Clotman, M.C. de Ruiter, A. Gittenberger-de Groot, R. Poelmann, F. Lupu, J.M. Herbert, D. Collen and E. Dejana: Targeted deficiency or cytosolic truncation of the VE-cadherin gene in mice impairs VEGF-mediated

Phosphatidylinositol 3-kinase

- endothelial survival and angiogenesis. *Cell* 98, 147-157 (1999)
72. J. Espada, M. Pérez-Moreno, V.M. Braga, P. Rodriguez-Viciana and A. Cano: H-Ras activation promotes cytoplasmic accumulation and phosphoinositide 3-OH kinase association of beta-catenin in epidermal keratinocytes. *J Cell Biol* 146, 967-980 (1999)
73. E. Calautti, J. Li, S. Saoncella, J.L. Brissette and P.F. Goetinck: Phosphoinositide 3-kinase signaling to Akt promotes keratinocyte differentiation versus death. *J Biol Chem* 280, 32856-32865 (2005)
74. L. Funke, S. Dakoji and D.S. Bredt: Membrane-associated guanylate kinases regulate adhesion and plasticity at cell junctions. *Annu Rev Biochem* 74, 219-245 (2005)
75. V.M. Braga: Cell-cell adhesion and signaling. *Curr Opin Cell Biol* 14, 546-556 (2002)
76. P. Humbert, S. Russell and H. Richardson: Dlg, Scribble and Lgl in cell polarity, cell proliferation and cancer. *Bioessays* 25, 542-553 (2003)
77. O. Bossinger, A. Klebes, C. Segbert, C. Theres and E. Knust: Zonula adherens formation in *Caenorhabditis elegans* requires dlg-1, the homologue of the *Drosophila* gene discs large. *Dev Biol* 230, 29-42 (2001)
78. L. McMahon, R. Legouis, J.L. Vonesch and M. Labouesse: Assembly of *C. elegans* apical junctions involves positioning and compaction by LET-413 and protein aggregation by the MAGUK protein DLG-1. *J Cell Sci* 114, 2265-2277 (2001)
79. B.L. Firestein and C. Rongo: DLG-1 is a MAGUK similar to SAP97 and is required for adherens junction formation. *Mol Biol Cell* 12, 3465-3475 (2001)
80. R.A. Lue, E. Brandin, E.P. Chan and D. Branton: Two independent domains of hDlg are sufficient for subcellular targeting: the PDZ1-2 conformational unit and an alternatively spliced domain. *J Cell Biol* 135, 1125-1137 (1996)
81. D.F. Woods, C. Hough, D. Peel, G. Callaini and P.J. Bryant: Dlg protein is required for junction structure, cell polarity, and proliferation control in *Drosophila* epithelia. *J Cell Biol* 134, 1469-1482 (1996)
82. D. Korkin, F.P. Davis, F. Alber, T. Luong, M.Y. Shen, V. Lucic, M.B. Kennedy and A. Sali: Structural modeling of protein interactions by analogy: application to PSD-95. *PLoS Comput Biol* 2, e153 (2006)
83. S.M. Reuver and C.C. Garner: E-cadherin mediated cell adhesion recruits SAP97 into the cortical cytoskeleton. *J Cell Sci* 111, 1071-1080 (1998)
84. H. Wu, S.M. Reuver, S. Kuhlendahl, W.J. Chung and C.C. Garner: Subcellular targeting and cytoskeletal attachment of SAP97 to the epithelial lateral membrane. *J Cell Sci* 111, 2365-2376 (1998)
85. P. Lévy, H. Robin, M. Kornprobst, J. Capeau and G. Cherqui: Enterocytic differentiation of the human Caco-2 cell line correlates with alterations in integrin signaling. *J Cell Physiol* 177, 618-627 (1998)
86. I. Royal and M. Park: Hepatocyte growth factor-induced scatter of Madin-Darby canine kidney cells requires phosphatidylinositol 3-kinase. *J Biol Chem* 270, 27780-27787 (1995)
87. S. Potempa and A.J. Ridley: Activation of both MAP kinase and phosphatidylinositide 3-kinase by Ras is required for hepatocyte growth factor/scatter factor-induced adherens junction disassembly. *Mol Biol Cell* 9, 2185-2200 (1998)
88. L.M. Shaw, I. Rabinovitz, H.H. Wang, A. Toker and A.M. Mercurio: Activation of phosphoinositide 3-OH kinase by the alpha6beta4 integrin promotes carcinoma invasion. *Cell* 91, 949-960 (1997)
89. F. Hollande, A. Choquet, E.M. Blanc, D.J. Lee, J.P. Bali and G.S. Baldwin: Involvement of phosphatidylinositol 3-kinase and mitogen-activated protein kinases in glycine-extended gastrin-induced dissociation and migration of gastric epithelial cells. *J Biol Chem* 276, 40402-40410 (2001)
90. S. Kermorgant, T. Aparicio, V. Dessirier, M.J. Lewin and T. Lehy: Hepatocyte growth factor induces colonic cancer cell invasiveness via enhanced motility and protease overproduction. Evidence for PI3 kinase and PKC involvement. *Carcinogenesis* 22, 1035-1042 (2001)
91. C. Bierkamp, A. Kowalski-Chauvel, S. Dehez, D. Fourmy, L. Pradayrol and C. Seva: Gastrin mediated cholecystokinin-2 receptor activation induces loss of cell adhesion and scattering in epithelial MDCK cells. *Oncogene* 21, 7656-7670 (2002)
92. K. Vleminckx and R. Kemler: Cadherins and tissue formation: integrating adhesion and signaling. *Bioessays* 21, 211-220 (1999)
93. K.M. Schmidt-Ott, D. Lan, B.J. Hirsh and J. Barasch: Dissecting stages of mesenchymal-to-epithelial conversion during kidney development. *Nephron Physiol* 104, 56-60 (2006)
94. S. Goossens and F. van Roy: Cadherin-mediated cell-cell adhesion in the testis. *Front Biosci* 10, 398-419 (2005)
95. C. Côme, V. Arnoux, F. Bibeau and P. Savagner: Roles of the transcription factors snail and slug during mammary morphogenesis and breast carcinoma progression. *J Mammary Gland Biol Neoplasia* 9, 183-193 (2004)

Phosphatidylinositol 3-kinase

96. K. Sundfeldt: Cell-cell adhesion in the normal ovary and ovarian tumors of epithelial origin; an exception to the rule. *Mol Cell Endocrinol* 202, 89-96 (2003)
97. A. Sue Menko: Lens epithelial cell differentiation. *Exp Eye Res* 75, 485-490 (2002)
98. U. Tepass, G. Tanentzapf, R. Ward and R. Fehon: Epithelial cell polarity and cell junctions in *Drosophila*. *Annu Rev Genet* 35, 747-784 (2001)
99. L. Larue, C. Antos, S. Butz, O. Huber, V. Delmas, M. Dominis and R. Kemler: A role for cadherins in tissue formation. *Development* 122, 3185-3194 (1996)
100. D. Riethmacher, V. Brinkmann and C. Birchmeier: A targeted mutation in the mouse E-cadherin gene results in defective preimplantation development. *Proc Natl Acad Sci USA* 92, 855-859 (1995)
101. M.L. Hermiston and J.I. Gordon: Inflammatory bowel disease and adenomas in mice expressing a dominant negative N-cadherin. *Science* 270, 1203-1207 (1995)
102. P. Young, O. Boussadia, H. Halfter, R. Grose, P. Berger, D.P. Leone, H. Robenek, P. Charnay, R. Kemler and U. Suter: E-cadherin controls adherens junctions in the epidermis and the renewal of hair follicles. *EMBO J* 22, 5723-5733 (2003)
103. C.L. Tinkle, T. Lechler, H.A. Pasolli and E. Fuchs: Conditional targeting of E-cadherin in skin: insights into hyperproliferative and degenerative responses. *Proc Natl Acad Sci USA* 101, 552-557 (2004)
104. A. Somasiri, C. Wu, T. Ellchuk, S. Turley and C.D. Roskelley: Phosphatidylinositol 3-kinase is required for adherens junction-dependent mammary epithelial cell spheroid formation. *Differentiation* 66, 116-125 (2000)
105. S.H. Kim, Z. Li and D.B. Sacks: E-cadherin-mediated cell-cell attachment activates Cdc42. *J Biol Chem* 275, 36999-37005 (2000)
106. M. Houde, P. Laprise, D. Jean, M. Blais, C. Asselin and N. Rivard: Intestinal epithelial cell differentiation involves activation of p38 mitogen-activated protein kinase that regulates the homeobox transcription factor CDX2. *J Biol Chem* 276, 21885-21894 (2001)
107. P. Laprise, M.J. Langlois, M.J. Boucher, C. Jobin and N. Rivard: Down-regulation of MEK/ERK signaling by E-cadherin-dependent PI3K/Akt pathway in differentiating intestinal epithelial cells. *J Cell Physiol* 199, 32-39 (2004)
108. J.C. Aliaga, C. Deschênes, J.F. Beaulieu, E.L. Calvo and N. Rivard: Requirement of the MAP kinase cascade for cell cycle progression and differentiation of human intestinal cells. *Am J Physiol* 277, G631-G641 (1999)
109. N. Rivard, M.J. Boucher, C. Asselin and G. L'Allemain: MAP kinase cascade is required for p27 down-regulation and S phase entry in fibroblasts and epithelial cells. *Am J Physiol* 277, C652-C664 (1999)
110. P.J. Van Haastert and P.N. Devreotes: Chemotaxis: signaling the way forward. *Nat Rev Mol Cell Biol* 5, 626-634 (2004)
111. S.J. Watton and J. Downward: Akt/PKB localisation and 3' phosphoinositide generation at sites of epithelial cell-matrix and cell-cell interaction. *Curr Biol* 9, 433-436 (1999)
112. A. Gassama-Diagne, W. Yu, M. ter Beest, F. Martin-Belmonte, A. Kierbel, J. Engel and K. Mostov: Phosphatidylinositol-3,4,5-trisphosphate regulates the formation of the basolateral plasma membrane in epithelial cells. *Nat Cell Biol* 8, 963-970 (2006)
113. F. Martin-Belmonte, A. Gassama, A. Datta, W. Yu, U. Rescher, V. Gerke and K. Mostov: PTEN-mediated apical segregation of phosphoinositides controls epithelial morphogenesis through Cdc42. *Cell* 128, 383-397 (2007)
114. K.D. Liu, A. Datta, W. Yu, P.R. Brakeman, T.S. Jou, M.A. Matthay and K.E. Mostov: Rac1 is required for reorientation of polarity and lumen formation through a PI 3-kinase-dependent pathway. *Am J Physiol Renal Physiol* 293, F1633-1640 (2007)
115. L.E. O'Brien, T.S. Jou, A.L. Pollack, Q. Zhang, S.H. Hansen, P. Yurchenco and K.E. Mostov: Rac1 orientates epithelial apical polarity through effects on basolateral laminin assembly. *Nat Cell Biol* 3, 831-838 (2001)

Abbreviations: PI3K: phosphatidylinositol 3-kinase; PtdIns: phosphoinositides; PH: pleckstrin homology; PKB: protein kinase B; MDCK: Madin-Darby Canine Kidney; SH domain: src homology domain; Dlg: disc large; PDZ: PSD-95/Dlg/ZO-1; MAGUK: membrane-associated guanylate cyclase; ERK: extracellular signal-regulated kinase; FAK: focal adhesion kinase

Key Words: Adherens junctions, phosphatidylinositol 3-kinase, PI3K, Phosphatidylinositol-3,4,5-trisphosphate, Phosphoinositides, Cell Adhesion, Epithelium, E-Cadherin, Actin Cytoskeleton, Catenin, Dlg, Review

Send correspondence to: Nathalie Rivard, Department of Anatomy and Cell Biology, Faculty of Medicine and Health Sciences, University of Sherbrooke, 3001 12th north avenue, Sherbrooke, Quebec, Canada, J1H5N4, Tel: 819-564-5271, Fax: 819-564-5320, E-mail: Nathalie.Rivard@USherbrooke.ca

<http://www.bioscience.org/current/vol14.htm>