

Paradoxical impact of cholesterol on lipid packing and cell stiffness

Manuela A Ayee¹, Irena Levitan¹

¹*Division of Pulmonary Care, Department of Medicine, University of Illinois at Chicago*

TABLE OF CONTENTS

1. Abstract
2. Introduction
3. Cholesterol-induced ordering of membrane phospholipids in artificial membranes
4. Impact of cholesterol on physical properties of plasma membrane lipid bilayers
 - 4.1. Experimental tools for modulating cellular cholesterol
 - 4.2. Cholesterol effects on the fluidity of plasma membranes
 - 4.3. Cholesterol effects on lipid packing of the plasma membrane bilayer
5. Impact of cholesterol on membrane-cytoskeleton interactions
 - 5.1. Impact of cholesterol on the lateral mobility of membrane proteins
 - 5.2. Cholesterol loss increases the strength of membrane-cytoskeleton adhesion
6. A loss of cholesterol results in cell stiffening
 - 6.1. Cell membrane deformability
 - 6.2. Stiffness of the intracellular “deep” cytoskeleton
 - 6.3. Force generation
7. Inverse effects of oxLDL and oxysterols on lipid packing and cell stiffness
 - 7.1. OxLDL-induced fluidization and stiffening of endothelial membranes
 - 7.2. 7-ketocholesterol-induced fluidization and stiffening of endothelial membranes
8. Concluding remarks and implications for cellular function
9. Acknowledgements
10. References

1. ABSTRACT

Cell stiffness or deformability is a fundamental property that is expected to play a major role in multiple cellular functions. It is well known that cell stiffness is dominated by the intracellular cytoskeleton that, together with the plasma membrane, forms a membrane-cytoskeleton envelope. However, our understanding of how lipid composition of plasma membrane regulates physical properties of the underlying cytoskeleton is only starting to emerge. In this review, we first briefly describe the impact of cholesterol on the physical properties of lipid bilayers in model membranes and in living cells, with the dominant effect of increasing the order of membrane lipids and decreasing membrane fluidity. Then, we discuss accumulating evidence that removal of cholesterol, paradoxically, decreases the mobility of membrane proteins and increases cellular stiffness, with both effects being dependent on the integrity of the cytoskeleton. Finally, we discuss emerging evidence that oxidized modifications of low-density lipoproteins (oxLDL) have the same effects on endothelial biomechanical properties as cholesterol depletion, an effect that is mediated by the incorporation of oxysterols into the membrane.

2. INTRODUCTION

Cholesterol is one of the major lipid components of the plasma membrane in all mammalian cells, where

it constitutes up to 45 mol% with respect to other lipids. Remarkably, there is a strong heterogeneity in cholesterol distribution between the plasma and the intracellular membranes, with the majority of cholesterol, up to 90%, found in the plasma membrane (1, 2). Furthermore, cholesterol is also distributed heterogeneously within the plasma membrane, giving rise to a concept of cholesterol-rich membrane rafts/domains (3, 4). The impact of cholesterol on the physical properties of the membrane lipid bilayer is well studied, both in pure lipid environments, such as liposomes or lipid monolayers, and in cellular membranes. Briefly, an increase in membrane cholesterol has been shown to increase lipid packing and decrease fluidity and deformability of lipid bilayers. However, an increasing number of studies, including by our group and other investigators, demonstrate that an increase in the rigidity and lipid packing of the bilayer does not translate into an increase in the overall stiffness of the cellular envelope, a bi-component system where the sub-membrane cytoskeleton underlies the membrane lipid bilayer. In contrast, it appears that there is an inverse relationship between the lipid order of the membrane bilayer and the stiffness of the cellular envelope that is dominated by the cortical cytoskeleton. The goal of this review is to discuss the current knowledge about the




Figure 1. Schematic of membrane thickening and phospholipid tail alignment caused by inserted cholesterol molecules. The phospholipid molecules (pink spheres with grey tails) form a bilayer. Cholesterol molecules are represented by red spheres for hydroxyl groups with yellow steroid rings and a black hydrophobic tail.

impact of membrane cholesterol on cellular biomechanics and the implications of these effects on cellular function.

3. CHOLESTEROL-INDUCED ORDERING OF MEMBRANE PHOSPHOLIPIDS IN ARTIFICIAL MEMBRANES

The cholesterol molecule consists of a planar ring structure built of four fused steroid rings with a hydroxyl group and a hydrophobic tail. The molecule is oriented with the steroid rings parallel to the hydrocarbon chains of the membrane phospholipids: the hydroxyl group interacts with the polar head groups of the phospholipids while the rigid lipophilic steroid ring system interacts with the hydrocarbon chains, restricting their motion within the bilayer (2, 5).

At low cholesterol concentrations (below 15 mol%), the lateral order of artificial phospholipid membranes made up of saturated phospholipids such as DPPC or DMPC is initially disrupted, causing membrane fluidization at room temperature, an effect that is reversed as cholesterol concentration increases (e.g. (6, 7)). Addition of cholesterol also abolishes the phase transition from the liquid to the gel phase of the model membrane (8). At the same time, however, above the phase transition temperature in membranes, the presence of cholesterol causes condensation of the phospholipids and a decrease in membrane fluidity. As discussed in detail below, the predominant effect of cholesterol in biological membranes is a decrease in membrane fluidity. The mechanism of the condensation effect is related to the energetic requirement of the membrane to shield the hydrophobic portions of cholesterol molecules from the external polar environment. For membrane phospholipids, their hydrophobic tails are shielded from the external polar environment by their large hydrophilic headgroups. Cholesterol, however, possesses a smaller hydrophilic headgroup than phospholipids; therefore, when present in a membrane, cholesterol is unable to shield the hydrophobic region as efficiently as phospholipids. To better shield the hydrophobic interior of the membrane in the presence of cholesterol, the phospholipid tails

become more aligned as they pack closer together, and the area they occupy decreases (9). Thus, an increase in membrane cholesterol concentration has an ordering effect on the membrane phospholipid tails, which also manifests itself as a decrease in membrane fluidity due to hindering of the lateral motion of membrane phospholipids, as described in earlier studies (e.g. (10-12)).

In terms of membrane elasticity, cholesterol has been shown to increase the elastic compressibility modulus of lipid bilayers, making the membrane effectively more rigid and less deformable (13, 14). Moreover, the heterogeneous distribution of membrane cholesterol, influenced by its differential interactions with phospholipids with varied degrees of acyl tail saturation, results in a non-monotonous change in the membrane elasticity modulus as cholesterol concentrations are increased (15-17). The decrease in in-plane membrane elasticity with increasing cholesterol concentration has been found to also depend on the degree of phospholipid acyl chain saturation and headgroup identity. Phospholipid membranes with greater tail saturation have greater decreases in elasticity when exposed to higher cholesterol concentrations (18, 19). This effect results in the formation of a liquid ordered phase in which phospholipid acyl tails appear to be extended and tightly packed, leading to the formation of membrane ordered domains (e.g. (20-22)). The ordering effect of cholesterol is also associated with increased thickness of the hydrophobic core of the membrane. For example, the thickness of saturated DPPC bilayers was shown to increase with increasing cholesterol concentration from 57.0 Å in bilayers without cholesterol, to 59.6 Å and 60.2 Å in bilayers containing 9% and 28% cholesterol respectively (23). This ordering effect is also attributed to an increase in *trans* configurations of the hydrocarbon chains in the case of unsaturated phospholipids. Accordingly, cholesterol-rich ordered membrane domains are expected to have a higher thickness than the rest of the membrane (Figure 1). Accumulating evidence suggests that cholesterol-rich membrane domains that are formed in cellular plasma membranes play an important role in the regulation of cellular stiffness.

4. IMPACT OF CHOLESTEROL ON PHYSICAL PROPERTIES OF PLASMA MEMBRANE LIPID BILAYERS

4.1. Experimental tools for modulating cellular cholesterol

A transition from studying effects of cholesterol on the properties of artificial membranes to cellular membranes presented an immediate challenge of modulating cell membrane cholesterol in a specific and reproducible way. Prior to the development of β -cyclodextrin (β CD)-based cholesterol extraction, the means to modify cellular cholesterol in a controlled way were limited. It was possible to incubate cells with liposomes containing either low or high cholesterol concentrations, which would respectively decrease or increase cellular cholesterol levels upon fusion to cell membranes. Alternatively, it was possible to serum starve the cells by maintaining them in media containing little to no serum. Clearly, both approaches introduce multiple non-specific effects. A major advance in the field was the development of a method to specifically modulate membrane cholesterol in cellular membranes in a precise and reproducible way by exposing cells to β -cyclodextrins (β CDs) (24, 25), which led to an explosion of studies addressing the role of cholesterol in a variety of cellular functions.

Cyclodextrins, oligosaccharides consisting of α -(1-4)-linked D-glycopyranose units, have long been utilized as carriers of hydrophobic drugs due to their high solubility in water and the presence of a hydrophobic cavity within the molecule (26). The size of the cavity depends on the degree of the polymerization and provides relative specificity for different hydrophobic loads (26), (27). Specifically, heptamers, or β -cyclodextrins, were shown to be the most efficient at extracting cholesterol, whereas hexamers, or α -cyclodextrins are more efficient at extracting phospholipids (28). In terms of extraction mechanism, Rothblatt and colleagues (29) proposed that, based on the kinetic and energetic analysis of cholesterol efflux, a cyclodextrin molecule diffuses into the immediate proximity of the plasma membrane bilayer and cholesterol molecules diffuse directly into the cyclodextrin hydrophobic cavity without having to dissolve into the aqueous phase of the cytosol.

It is important to note that, while numerous studies established that β -cyclodextrins can be used for precise and reproducible cholesterol extraction, enrichment or replenishment in multiple cell types and tissues, the specificity of cholesterol extraction for cholesterol-rich vs. cholesterol-poor membrane domains remains controversial. Briefly, some specificity was suggested by studies showing that short exposures (≤ 2 min) or very low concentrations (≤ 1 mM) of M β CD result in selective extraction of cholesterol from cholesterol-rich membrane domains (30, 31), however, the preponderance of studies

show that M β CD is capable of removing cholesterol from both types of membrane domains (27). A more detailed analysis of the impact of cholesterol depletion and enrichment on membrane fluidity and local lipid order of plasma membranes is presented below.

4.2. Cholesterol effects on the fluidity of plasma membranes

Membrane fluidity, an inverse of viscosity, can be estimated by measuring the motion of different moieties within the bilayer. The two most common methods to measure membrane fluidity are *fluorescence anisotropy*, based on the rotational diffusion of fluorescent lipid probes, and *FRAP* (Fluorescence Recovery After Photobleaching), an approach that measures the rate of lateral diffusion of a fluorescent moiety into a region that was previously photobleached.

Fluorescence anisotropy is based on the incorporation of small fluorescent lipophilic probes into the membrane bilayer that orient themselves with the hydrocarbon chains of the phospholipids. Membrane viscosity is estimated from the rotational diffusion coefficients of the probes, which is hindered by the neighboring lipids (32-34). The probes differ in their selectivity for plasma vs. intracellular membranes and also in their specific position within the bilayer. For example, DPH (1,6-diphenyl-1,3,5-hexatriene) is incorporated into the hydrophobic core of the bilayer and has poor selectivity between the plasma and intracellular membranes, whereas its derivative TMA-DPH (4-trimethyl-ammonio-1,6-diphenyl-1,3,5-hexatriene) is incorporated in the outer leaflet close to the membrane surface and is more selective to the plasma membrane (32, 35). Universally, the effect of cholesterol removal is an increase in membrane fluidity, as measured by DPH-based fluorescence anisotropy. Specific examples include a significant decrease in fluorescence anisotropy corresponding to increased membrane fluidity in cholesterol depleted HEK293 cells (36, 37), membrane vesicles isolated from a basophilic leukemia cell line (38), human glioblastoma cells (39), primary cerebellar granule cells (40) and sperm (41).

Effects of cholesterol depletion on the lateral mobility of lipid probes or lipid components of biological membranes are mostly consistent with increased membrane fluidity reported by fluorescence anisotropy. Pucadyil and Chattopadhyay (42) reported that cholesterol depletion of hippocampal membranes resulted in a significant increase in lateral diffusion of two fluorescent lipids: DiI_{C₁₈}, which partitions preferentially into ordered domains, and FAST DiI, which partitions into disordered membranes. Interestingly, the effect of cholesterol depletion on lipid diffusion was higher for ordered than for disordered phases. Cholesterol depletion was also reported to increase the lateral mobility of a fluorescent analogue of sphingomyelin (43). A recent study showed,




Figure 2. Impact of cholesterol depletion on lipid packing of membrane domains in endothelial cells. A: Typical GP images of control and cholesterol-depleted cells. Bar is 5.6 μm. B: The zoomed-in representative regions of the GP images shown above. Scale bar is 1 μm. All images are shown in pseudo color with yellow and red corresponding to the higher GP values, presumably ordered domains, and green and blue corresponding to the lower GP values, presumably disordered domains. C: GP histograms fitted by a two-Gaussian distribution. Reproduced from Shentu *et al.*, 2010, American J of Physiology Cell Physiology 299: C218-229.

however, that cholesterol depletion may also decrease the mobility of a lipid probe in primary hair cells isolated from guinea pigs, an effect that was specific to the basal membranes, while no effect was observed on the apical membranes (44). The source of this local heterogeneity is not clear. In any case, an increase in membrane fluidity is expected to be associated with increased lipid packing/ordering of the membrane.

4.3. Cholesterol effects on lipid packing of the plasma membrane bilayer

Lipid packing or ordering of biological membranes can be estimated by Laurdan two-photon microscopy, a dye that is sensitive to the polarity of the local environment and undergoes a red shift as the phase boundary changes from gel to fluid (45, 46). Changes in

membrane order are estimated by calculating the general polarization (GP) ratio, a normalized ratio of fluorescence intensity of gel phase vs. fluid phase. Typically, plasma membranes are shown to be highly heterogeneous in their lipid order with GP values varying in a continuous manner and without a sharp, clear distinction between “ordered” and “disordered” domains. Nevertheless, GP histograms reflect the relative abundance of the membrane domains with more fluid or more ordered properties, providing a “map” of lipid order of individual cells. Figure 2 shows a typical example of a “domain map”, as detected by Laurdan two-photon microscopy.

Gaus *et al.* (45) were first to test the effects of cholesterol depletion and enrichment on the lipid order of membrane domains using Laurdan imaging. Their data

show that cholesterol depletion of macrophages results in a significant decrease in lipid order of both ordered and fluid domains, and a relatively small, but significant, shift in the total area of the cell membrane covered by ordered vs. fluid domains (about a 10-20% shift from ordered to fluid). Cholesterol enrichment, however, did not increase the order of either of the domains, but increased the area covered by the ordered domains (also by about 20%). A decrease in lipid order following cholesterol depletion was also observed in plasma membranes of multiple other cell types, including kidney cells (47), hippocampal membranes (48), and aortic endothelial cells (49), as well as in cell lines CHO (50) and HEK (37). Interestingly, M β CD-induced cholesterol depletion was observed not only to decrease lipid order, but also to decrease membrane heterogeneity (49).

In summary, it is well established that removal of cellular cholesterol increases membrane fluidity, as measured by the diffusion of lipid probes, and decreases the lipid order/packing of the membrane lipid bilayer of plasma membranes. This effect is based on the physical interactions between cholesterol molecules and phospholipids, and is fully consistent with the impact of cholesterol on fluidity and lipid packing of non-biological membranes.

An increase in membrane fluidity should be expected to promote the ability of membrane proteins to move within the membrane and form regulatory protein-protein complexes, which clearly should have major functional consequences. However, accumulating evidence demonstrates that cholesterol depletion actually inhibits the lateral motilities of membrane proteins in several systems, an effect that depends on the integrity of the sub-membrane cytoskeleton.

5. IMPACT OF CHOLESTEROL ON MEMBRANE-CYTOSKELETON INTERACTIONS

5.1. Impact of cholesterol on the lateral mobility of membrane proteins

The first surprising result came from the work of Edidin and colleagues, who used FRAP to determine the effect of cholesterol depletion on the lateral diffusion of a class of HLA (Human Leukocyte Antigen) molecules in fibroblasts and lymphoblasts (51). In contrast to an increase in membrane fluidity reported by fluorescence anisotropy of membrane probes, Kwik *et al.* showed that cholesterol depletion significantly decreased the mobility of the HLA molecules. A similar result was also observed for epidermal growth factor receptors in the same cells. A decrease in the lateral mobility of the membrane proteins in cholesterol-depleted cells was abrogated by depolymerizing F-actin, indicating that this effect depends on the stability of the cytoskeleton (51). A similar effect was observed for voltage-gated K⁺ channels Kv1.4., with

cholesterol depletion resulting in a significant decrease in the lateral mobility of the channels (52). At the same time, the mobilities of two other voltage-gated K⁺ channels, Kv2.1. and Kv1.3., were either unaffected or slightly increased, leading to the conclusion that these channels partition into separate membrane domains. The exact mechanisms of the differential effects of cholesterol on Kv channels, however, remain unclear. A decrease in mobility following cholesterol depletion was also reported for several other proteins, including Serotonin_{1A} receptors (53) and Nicotinic acetylcholine receptor (54). The major effects were a decrease in the mobile fraction and of the diffusion coefficients. Interestingly, Baier *et al.* also showed that depolymerization of F-actin resulted in partial recovery of receptor mobility in cholesterol-depleted cells (54). Taken together, these observations lead to two conclusions: lateral mobility of membrane proteins may be inversely related to the fluidity of lipid components of the membrane, and the mobility is critically dependent on the integrity of the cytoskeleton. These observations also imply that, surprisingly, the integrity/stability of the sub-membrane cytoskeleton and/or membrane-cytoskeleton adhesion is enhanced by cholesterol depletion. Direct evidence for this effect is discussed in the next section.

5.2. Cholesterol loss increases the strength of membrane-cytoskeleton adhesion

Earlier studies investigated the role of cholesterol-rich membrane domains in membrane-cytoskeleton interactions with accumulating evidence suggesting that these domains serve as focal points for these interactions. More specifically, several major regulators of the cytoskeleton function, such as phosphatidylinositol 4,5 bisphosphate (PIP₂) and Rho-GTPases were found to partition into the protein complexes found in raft domains (55). Cholesterol depletion, on the other hand, was shown to disrupt the association of PIP₂ with the raft domains (56), decrease the level of PIP₂ on the plasma membrane (51), and prevent membrane association and activation of Rac1 (57). These studies were performed in several cell types including fibroblasts and lymphoblasts, as well as in immortalized cell lines. Furthermore, multiple cytoskeleton proteins were identified in cholesterol-rich detergent-resistant membrane fractions in neutrophils using proteomic analysis (58). Based on these studies, the expectation was that cholesterol depletion resulting in the disruption of cholesterol-rich membrane domains should result in weakening of membrane-cytoskeleton interactions and dissociation of the sub-membrane cytoskeleton from the plasma membrane. Thus, studies of the effect of cholesterol depletion on the lateral mobility of membrane proteins and studies of the association of cytoskeleton proteins with cholesterol-rich domains created opposite expectations regarding the effect of membrane cholesterol on the strength of membrane-cytoskeletal adhesion. It was critical, therefore, to determine directly how changes in the level of membrane

cholesterol regulate the adhesion between the plasma membrane and the sub-membrane cytoskeleton.

Sun *et al.* (59) investigated the impact of cholesterol on the adhesion strength between the cytoskeleton and the membrane by pulling membrane tethers from the surface of aortic endothelial cells using Atomic Force Microscopy (AFM). This approach, described in detail by (60), measures the tether force (F), recorded by the deformation of the AFM cantilever, as the cantilever is pulled away from the cells at different speeds. The dependence of the tether force and the pulling speed is described by $F = F_0 + 2\pi\eta_{eff}v$ (61), where F_0 , the threshold force, reflects the amount of force needed to initiate a tether formation and is a function of the membrane-cytoskeleton interaction, η_{eff} is the effective membrane surface viscosity, and v is the speed at which the cantilever is pulled away from the cell. Using this approach, we found that cholesterol depletion significantly increased the threshold force that is required to pull the tether, which is directly related to the membrane-cytoskeleton interaction (59). An increase F_0 in was abrogated by depolymerization of F-actin, further suggesting that this effect should be attributed to the stabilization and strengthening of the sub-membrane cytoskeleton. To isolate the contribution of actin-cytoskeleton in determining the F_0 of generating the tether, we defined the membrane-cytoskeleton adhesion, F_{ad} as the difference in F_0 before and after latrunculin treatment, i.e. $F_{ad} = F_0 - F_0^{lata}$. Cholesterol depletion results in almost a 2-fold increase in F_{ad} , whereas cholesterol enrichment significantly decreases the threshold tether force F_0 so that it becomes almost equivalent to F_0^{lata} . These observations led to the conclusion that cholesterol depletion increases membrane-cytoskeleton adhesion, whereas cholesterol enrichment has the opposite effect. Furthermore, cholesterol depletion also increases the variance of tether force, suggesting that cholesterol depletion makes tethers much more heterogeneous, presumably due to the enhanced membrane-cytoskeleton adhesion. In contrast, cholesterol enrichment reduces the membrane-cytoskeleton adhesion and makes the bilayer easier to detach from and flow on the cytoskeleton.

In terms of the mechanism, our more recent study suggests that cholesterol depletion enhances membrane-cytoskeletal interactions by causing dispersion and redistribution of PIP₂ (62). Similarly to the previous study, we used aortic endothelial cells to show that while PIP₂ appeared in discrete patches in control endothelial cells, cholesterol depletion resulted in the loss of discrete PIP₂ domains and almost uniform distribution across the cell membrane. Most importantly, while addition of exogenous PIP₂ had no effect on the tether force in control cells, it significantly increased the tether force in cholesterol-depleted cells indicating stronger membrane-cytoskeleton adhesion. We hypothesized that cholesterol depletion may strengthen membrane-cytoskeleton interactions by increasing the number

of focal points enriched with the PIP₂ linker. We could not completely exclude the possibility, however, that cholesterol depletion makes it easier for the exogenous PIP₂ to be incorporated into the plasma membrane, thus more studies are needed to evaluate this point. Interestingly, changes in PIP₂ distribution were also shown to be critical for the effect of cholesterol depletion on constraining the lateral mobility of membrane proteins reported earlier (51), but in that case, cholesterol depletion resulted in the loss of PIP₂ from the plasma membrane. The authors suggested that the loss of PIP₂ results in re-organization and stabilization of the sub-membrane cytoskeleton. Clearly, more studies are needed to provide better understanding of these mechanisms.

6. A LOSS OF CHOLESTEROL RESULTS IN CELL STIFFENING

6.1. Cell membrane deformability

As described above, the first indication that cholesterol removal may result in the stabilization of the cytoskeleton came from a study by Kwik *et al.* (51) showing that cholesterol depletion of human fibroblasts constrained the mobility of a class of transmembrane proteins, an effect that was fully dependent on the integrity of F-actin. Furthermore, Kwik *et al.* proposed that cholesterol depletion resulted in stabilization of the sub-membrane cytoskeleton. Since the overall stiffness of the membrane-cytoskeleton complex is well recognized to be dominated by the rigidity of the sub-membrane cortical cytoskeleton (63), these observations implied that a loss of membrane cholesterol may have significant implications for the biomechanical properties of the membrane-cytoskeleton complex.

Byfield *et al.* (64) provided the first direct evidence that cholesterol depletion results in significant stiffening and loss of deformability of vascular endothelial cells by measuring progressive membrane deformation in response to negative pressure applied by a glass micropipette (Figure 3). Exactly the same approach was used earlier to determine the effect of cholesterol on membrane stiffness of liposomes, showing that an increase in membrane cholesterol results in bilayer stiffening (13). However, in contrast to its effect on the bilayer stiffness, removal of membrane cholesterol either by a cholesterol-depleting agent, methyl- β -cyclodextrin (M β CD), or by serum starvation resulted in the slower rate of membrane deformation and smaller maximal deformation in response to the same level of negative pressure, indicating an increase in endothelial stiffness (64). Cholesterol depletion-induced endothelial stiffening was accompanied with a significant change in endothelial morphology and spreading (65). Enriching the cells with cholesterol had no effect. Importantly, a typical membrane area that is aspirated into the pipette in these experiments is rather large (the tips of the pipettes are ~5 μ m in diameter) and the membrane is clearly




Figure 3. Cholesterol depletion increases stiffness of vascular endothelial cells. A: Typical images of membrane deformation in response to negative pressure in cells depleted or enriched with cholesterol as compared to a control cell. Membrane is visualized with a fluorescent dye Di_{18} ; bar is 10 μ m. B: Maximal membrane deformation normalized to a pipette diameter as a function of applied pressure. Reproduced from Byfield *et al.*, 2004, *Biophys J.* 2004 Dec;87(6):3850-61.

undergoing deformation together with at least some of the sub-membrane cytoskeleton. The stiffening effect induced by cholesterol depletion is fully abrogated by the depolymerization of F-actin. The latter observation is consistent with the notion that integrity of F-actin plays the dominant role in determining cellular biomechanical properties. Most of these studies were performed using bovine aortic endothelial cells as a cellular model. Importantly, an increase in cell stiffness upon cholesterol depletion was also shown in Chinese Hamster Ovary cells (CHO K1) demonstrating that this effect is not unique to endothelial cells (66). Clearly, more studies are needed to determine the relationship between membrane cholesterol and cellular stiffness in other vascular tissues, particularly in vascular smooth muscle cells.

6.2. Stiffness of the intracellular “deep” cytoskeleton

Another rather unexpected observation came from the analysis of correlated motion of fluorescent beads that were phagocytosed by endothelial cells (67). The motion of internalized particles is indicative of the material properties of a cellular milieu, with greater motion indicating a softer material (68). Furthermore, movement of individual particles deforms the cytoskeletal network around them, resulting in displacement of other particles in their vicinity (69). Thus, analysis of the correlated motion of particles within a cell allows estimation of the stiffness of the intracellular cytoskeleton (70, 71). Using this approach, we showed that cholesterol depletion results in a significant (~50%) decrease in the correlated

motion of particles within endothelial cells, indicating an increase in the intracellular stiffness. It is also important to note that a decrease in the correlated motion of the particles may also be indicative of a decrease in motor activity. In either case, these observations indicate that cholesterol depletion regulates the material properties of the intracellular cytoskeleton.

6.3. Force generation

An increase in cellular stiffness in cholesterol depleted endothelial cells correlates with an increase in endothelial force generation. First, it was established that decreasing the level of cellular cholesterol facilitates the ability of aortic endothelial cells to induce gel contraction (72), which is indicative of an increase in force generation on the cell-substrate interface. Indeed, these observations are consistent with earlier studies that demonstrated that increased cell stiffness correlates with the magnitude of forces that cells exert on substrates (73). To gain further insights into the effect of cholesterol loss on endothelial force generation, we analyzed the local forces generated by endothelial cells using Traction Force Microscopy (TFM), as described previously (74). In this study, we showed that consistent with the gel contraction assays, TFM analysis revealed that cholesterol depletion results in an almost 2-fold increase in force exerted by the cells on their substrates (75). Similar to the impact of cholesterol on endothelial stiffness, while cholesterol depletion increased endothelial force generation, cholesterol enrichment had no effect. Force generation effects of cholesterol depletion, however, can be rescued

by replenishing cellular cholesterol. Thus, it appears that for both endothelial stiffness and force generation, there is a critical cholesterol level that is required to maintain the normal parameters, so that addition of cholesterol over this level has no further effect.

7. INVERSE EFFECTS OF OXLDL AND OXYSTEROLS ON LIPID PACKING AND CELL STIFFNESS

With an increased realization that the loss of membrane cholesterol has opposite effects on the lipid order of the membrane vs. on membrane-cytoskeleton interactions and cellular stiffness, the question arises of how this effect is related to physiological and pathological conditions in dyslipidemia. The main concern is usually an increase in cholesterol load rather than cholesterol deficiency. Our studies, however, provide multiple lines of evidence that exposing endothelial cells to oxidized low-density lipoproteins (oxLDL) results in the same effects on endothelial biomechanics as cholesterol depletion. Specifically, these effects include disruption of lipid packing (49), increase in cell stiffness (49, 72), increase in force generation (72, 76), and enhanced sensitivity to shear stress (77). Furthermore, we also showed that exposing cells to 7-ketocholesterol, one of the major components of oxLDL, also disrupts lipid order and increases endothelial stiffening (78). All of these studies focused on aortic endothelial cells (bovine and human) because of the high prevalence of atherosclerosis in these arteries. It would be very interesting and important to compare the biomechanical properties of endothelial cells across different vascular beds. A more detailed discussion of the current studies is provided below.

7.1. OxLDL-induced fluidization and stiffening of endothelial membranes

Oxidative damage of low-density lipoproteins (LDL) is well known to be a major risk factor for the development of cardiovascular disease. The levels of oxLDL increase in hypercholesterolemia in both humans and animal models, and it is found in atherosclerotic lesions (79-81). It is important to note, however, that the term oxidized LDL is used to describe a wide variety of LDL modifications ranging from minimally or mildly-oxidized to strongly-oxidized, which may differ significantly in their composition and biological effects. This variability can be attributed to the fact that LDL oxidation *in vivo* is mediated by a variety of free radicals including superoxide, hydroxyl radicals and nitric oxide produced by enzymatic action of several enzymes, such as NADPH oxidase (NOX), NO synthase and lipoxygenase (82). The most frequently used oxidation route *in vitro* is exposure to free transition metals, particularly copper, although this pathway is not significant in the physiological oxidation of LDL *in vivo*. The rationale to use copper-oxidized LDL as a molecular model for oxidative modifications of LDL *in vivo* is that it is recognized by the cellular scavenger receptors and

results in significant endothelial dysfunction. One of the major postulates in the field regarding oxLDL has been that oxLDL is detrimental to endothelial function because by virtue of being recognized by the scavenger receptors, it should be more efficient in loading cells with cholesterol. This notion was challenged by the studies of Smart and colleagues, who showed that rather than enriching endothelial cells with cholesterol, oxLDL depleted cholesterol from caveolae (83). Following this study, we tested the impact of oxLDL on the deformability of aortic endothelial cells and found that, similarly to cholesterol depletion, exposure to oxLDL resulted in a significant increase in endothelial stiffness (72). The impact of oxLDL on endothelial stiffening correlated positively with the degree of LDL oxidation (78). These studies were conducted using copper-oxidized LDL. More recently, we have also determined that exposing endothelial cells to enzymatically oxidized LDL also results in endothelial stiffening (manuscript in preparation). Furthermore, the effect of oxLDL on endothelial stiffness was verified in endothelial cells freshly-isolated from hypercholesterolemic pigs demonstrating that it is highly physiologically relevant (72).

OxLDL-induced endothelial stiffening was accompanied with a decrease in lipid order/fluidization of endothelial membranes, as determined by Laurdan imaging described above (49). Notably, both oxLDL-induced fluidization of the membrane domains and the stiffening of the cellular envelope could be reversed by increasing cholesterol content of the membrane. Thus, both of these effects are clearly not mediated by an increase in membrane cholesterol. However, since we also did not detect any significant decrease in cholesterol content, efflux or internalization in oxLDL-treated cells, we proposed an alternative hypothesis that oxLDL-induced fluidization of the membrane and endothelial stiffening are mediated by the membrane insertion of oxysterols.

7.2. 7-ketocholesterol-induced fluidization and stiffening of endothelial membranes

OxLDL contains an array of oxidative lipids that are bioactive under pathological conditions. Our studies showed that oxLDL-induced endothelial stiffening can be accounted for by the oxysterol fraction of the lipoprotein, with the most prominent effects caused by 7-ketocholesterol. We also found increased incorporation of 7-ketocholesterol in arterial tissues of dyslipidemic mice (78). Figure 4 shows a shift in the distribution of endothelial elastic moduli to higher values in cells exposed to 7-ketocholesterol, indicating an increase in cell stiffness. At the same time, the lipid order parameters of endothelial membranes shift to a more fluid, less ordered phase, as indicated by a relative decrease in highly-ordered (yellow) domains and an increase in less ordered (blue-green) domains. In addition, 27-hydroxycholesterol, which was also




Figure 4. 7-ketocholesterol has inverse effects on lipid order and endothelial stiffness. A: Typical GC chromatographs of oxysterols in lipid extracts isolated from control cells; cells exposed to 10 g/ml 7-ketocholesterol. B: Histograms of elastic modulus measured in control (Ctrl) and 7-ketocholesterol-treated cells. C: Average elastic modulus for control (Ctrl) and 7-ketocholesterol-treated cells. (mean \pm SEM, n=80 cells for each experimental condition). D: Typical GP images of control cells (Ctrl), 7-ketocholesterol-treated cells. Scale bar is 11.2 μ m.




Figure 5. Schematic of membrane fluidization and stabilization of sub-membrane cytoskeleton following cholesterol depletion and oxysterol insertion. Coloring follows that of Figure 1, with the addition of oxysterols containing extra polar groups represented by green spheres. Sub-membrane cytoskeleton filaments are represented by violet strands.

found in atherosclerotic lesions, had a similar effect. The stiffening effects induced by both 7-ketocholesterol and 27-hydroxycholesterol were fully reversed by enriching the cells with cholesterol (78). In terms of the lipid order effect, similarly to oxLDL, exposure to 7-ketocholesterol also resulted in a significant shift of endothelial membranes to a more fluid/less ordered

state. The latter is consistent with earlier studies showing that 7-ketocholesterol disrupts lipid order of model membranes (84). It is important to note that an inverse relationship between fluidization of the membrane and endothelial stiffening was also observed for another sterol, coprostanol (49), indicating this effect is not unique to the oxysterol components of oxLDL.

8. CONCLUDING REMARKS AND IMPLICATIONS FOR CELLULAR FUNCTION

In conclusion, multiple lines of evidence demonstrate that removal of membrane cholesterol or incorporation of specific oxysterols into the plasma membrane result in fluidization of the membrane lipid bilayer, and at the same time, lead to the strengthening of membrane-cytoskeleton interactions and increase in cell stiffness. Furthermore, membrane incorporation of oxysterols appears to be responsible for endothelial stiffening when exposed to oxLDL. The mechanisms, however, that couple fluidization of the membrane to increases in cell stiffness are still poorly understood. One mechanism that contributes to strengthening of membrane-cytoskeleton adhesion and stabilizing of the sub-membrane cytoskeleton is a loss or re-distribution of the regulatory phospholipid, PIP₂ (51, 62). A schematic summary is shown in Figure 5. Other possible mechanisms may include activation of Rho-GTPases or other signaling molecules that regulate the stability of the cytoskeleton. Further studies are needed to unravel these mechanisms.

In terms of functional significance, an increase in cell stiffness is likely to affect a plethora of cellular functions. More specifically, an increase in endothelial stiffness was shown to impair generation of nitric oxide in response to shear stress forces generated by blood flow (85), but also to facilitate flow-induced realignment of endothelial cells, a hallmark of their sensitivity to flow (77) and enhance transmigration of neutrophils through the endothelial layer (76). A series of studies also implicate endothelial stiffness as an important factor in angiogenesis (49, 73, 86, 87). Furthermore, an increase in smooth muscle stiffness has been proposed to underlie the stiffening of blood vessels in hypertension and aging (88, 89). Thus, changes in cellular cholesterol/oxysterol compositions may have profound effects on all of these properties. Functional significance should also be further assessed by exploring the impact of different modifications of oxLDL and particularly glycation of LDL that occurs in diabetes promoting further oxidation of LDL and formation of other oxidized lipids. The relationship between the lipid composition of the membrane and cellular stiffness can be an important factor in multiple pathological conditions.

9. ACKNOWLEDGEMENTS

We are very grateful to Dr. Zhongkui Hong for the critical reading of this manuscript. We also thank Mr. Gregory Kowalsky for his help in the layout and formatting of the Figures and Ms. Elizabeth LeMaster for proofreading and comments. The work was supported by National Institutes of Health grants HL073965 and HL083298 (to I. Levitan).

10. REFERENCES

1. P. L. Yeagle: Modulation of membrane function by cholesterol. *Biochimie*, 73, 1303 (1991)
DOI: 10.1016/0300-9084(91)90093-G
2. P. L. Yeagle: Cholesterol and the cell membrane. *Biochim. Biophys. Acta*, 822, 267-287 (1985)
DOI: 10.1016/0304-4157(85)90011-5
3. K. Simons and D. Toomre: Lipid rafts and signal transduction. *Nature reviews. Molecular cell biology.*, 1, 31-9 (2000)
DOI: 10.1038/35036052
4. M. Edidin: The state of lipid rafts: From model membranes to cells. *Annual Review Biophysical Biomolecular Structure.*, 32, 257083 (2003)
DOI: 10.1146/annurev.biophys.32.110601.142439
5. H. Ohvo-Rekilä, B. Ramstedt, P. Leppimäki and J. P. Slotte: Cholesterol interactions with phospholipids in membranes. *Prog Lipid Res* 41, 66-97 (2002)
DOI: 10.1016/S0163-7827(01)00020-0
6. D. Needham, T. J. McIntosh and E. Evans: Thermomechanical and transition properties of dimyristoylphosphatidylcholine/cholesterol bilayers. *Biochemistry*, 27, 4668-73 (1988)
DOI: 10.1021/bi00413a013
7. K. J. Tierney, D. E. Block and M. L. Longo: Elasticity and Phase Behavior of DPPC Membrane Modulated by Cholesterol, Ergosterol, and Ethanol. *Biophys. J.*, 89(4), 2481 (2005)
DOI: 10.1529/biophysj.104.057943
8. M. F. N. Rosser, H. M. Lu and P. Dea: Effects of alcohols on lipid bilayers with and without cholesterol: the dipalmitoylphosphatidylcholine system. *Biophysical Chemistry*, 81(1), 33 (1999)
DOI: 10.1016/S0301-4622(99)00081-2
9. F. de Meyer and B. Smit: Effect of cholesterol on the structure of a phospholipid bilayer. *Proc Natl Acad Sci U S A*, 106, 3654-3658 (2009)
DOI: 10.1073/pnas.0809959106
10. R. A. Demel, K. R. Bruckdorfer and L. L. M. van Deenen: The effect of sterol structure on the permeability of liposomes to glucose, glycerol and Rb⁺. *Biochim. Biophys. Acta*, 255, 321-330 (1972)

- DOI: 10.1016/0005-2736(72)90031-4
11. R. A. Demel and B. De Kruffy: The function of sterols in membranes. *Biochim Biophys Acta*, 457, 109-132 (1976)
DOI: 10.1016/0304-4157(76)90008-3
 12. B. W. Stockton and I. C. P. Smith: A deuterium NMR study of the condensing effect of cholesterol on egg phosphatidylcholine bilayer membranes. *Chem Phys Lipids*, 17, 251-263 (1976)
DOI: 10.1016/0009-3084(76)90070-0
 13. D. Needham and R. S. Nunn: Elastic deformation and failure of lipid bilayer membranes containing cholesterol. *Biophys J.*, 58, 997-1009 (1990)
DOI: 10.1016/S0006-3495(90)82444-9
 14. E. Evans and D. Needham: Physical properties of surfactant bilayer membranes: thermal transition, elasticity, rigidity, cohesion and colloidal interactions. *J.I of Physical Chem.*, 91, 4219-28 (1987)
DOI: 10.1021/j100300a003
 15. T. Hianik and M. Haburcak: Clustering of cholesterol in DMPC bilayers as indicated by membrane mechanical properties. *Gen Physiol Biophys.*, 12, 283-91 (1993)
No DOI Found
 16. T. Hianik, M. Haburcak, K. Lohnerb, E. Prennerc, F. Paltaufc and A. Hermetterc: Compressibility and density of lipid bilayers composed of polyunsaturated phospholipids and cholesterol. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 139, 189-197 (1998)
DOI: 10.1016/S0927-7757(98)00280-5
 17. J. S. Pan, Tristram-Nagle and J. F. Nagle: Effect of cholesterol on structural and mechanical properties of membranes depends on lipid chain saturation. *Physical review. E, Statistical, nonlinear, and soft matter physics*, 80, 021931 (2009)
DOI: 10.1103/PhysRevE.80.021931
 18. J. M. Smaby, M. M. Momsen, H. L. Brockman and R. E. Brown: Phosphatidylcholine acyl unsaturation modulates the decrease in interfacial elasticity induced by cholesterol. *Biophys. J.*, 73(3), 1492 (1997)
DOI: 10.1016/S0006-3495(97)78181-5
 19. X.-M. Li, M. M. Momsen, J. M. Smaby, H. L. Brockman and R. E. Brown: Cholesterol Decreases the Interfacial Elasticity and Detergent Solubility of Sphingomyelins. *Biochemistry*, 40(20), 5954 (2001)
DOI: 10.1021/bi002791n
 20. J. H. Ipsen, O. G. Mouritsen and M. Bloom: Relationships between lipid membrane area, hydrophobic thickness, and acyl-chain orientational order. The effects of cholesterol. *Biophys. J.*, 57(3), 405 (1990)
DOI: 10.1016/S0006-3495(90)82557-1
 21. M. B. Sankaram and T. E. Thompson: Interaction of cholesterol with various glycerophospholipids and sphingomyelin. *Biochemistry*, 29, 10670-5 (1990)
DOI: 10.1021/bi00499a014
 22. C. Reyes Mateo, A. Ulises AcuÃ±a and J. C. Brochon: Liquid-crystalline phases of cholesterol/lipid bilayers as revealed by the fluorescence of trans-parinaric acid. *Biophys. J.*, 68(3), 978 (1995)
DOI: 10.1016/S0006-3495(95)80273-0
 23. E. Drolle, N. KuÅ erka, M. I. Hoopes, Y. Choi, J. Katsaras, M. Karttunen and Z. Leonenko: Effect of melatonin and cholesterol on the structure of DOPC and DPPC membranes. *Biochim. et Biophys. Acta - Biomembranes*, 1828(9), 2247 (2013)
DOI: 10.1016/j.bbamem.2013.05.015
 24. U. Klein, G. Gimple and F. Fahrenholz: Alteration of the myometrial plasma membrane cholesterol with b-cyclodextrin modulates the binding affinity of the oxytocin receptor. *Biochemistry*, 34, 13784-93 (1995)
DOI: 10.1021/bi00042a009
 25. A. E. Christian, M. P. Haynes, M. C. Phillips and G. H. Rothblat: Use of cyclodextrins for manipulating cellular cholesterol content. *J. Lipid Res.*, 38, 2264-2272 (1997)
No DOI Found
 26. K. Uekama: Design and evaluation of cyclodextrin-based drug formulation. *Chem Pharm Bull (Tokyo)*, 52(8), 900-15 (2004)
DOI: 10.1248/cpb.52.900
 27. R. Zidovetzki and I. Levitan: Use of cyclodextrins to manipulate plasma membrane cholesterol content: Evidence, misconceptions and control strategies. *Biochim. et Biophys. Acta - Biomembranes*, 1768(6), 1311 (2007)
DOI: 10.1016/j.bbamem.2007.03.026

28. Y. Ohtani, T. Irie, K. Uekama, K. Fukunaga and J. Pitha: Differential effects of alpha-, beta- and gamma-cyclodextrins on human erythrocytes. *Eur J Biochem*, 186(1-2), 17-22 (1989)
DOI: 10.1111/j.1432-1033.1989.tb15171.x
29. P. G. Yancey, W. V. Rodriguez, E. P. Kilsdonk, G. W. Stoudt, W. J. Johnson, M. C. Phillips and G. H. Rothblat: Cellular cholesterol efflux mediated by cyclodextrins. Demonstration of kinetic pools and mechanism of efflux. *J Biol Chem*, 271(27), 16026-34 (1996)
DOI: 10.1074/jbc.271.27.16026
30. A. K. Rouquette-Jazdanian, C. Pelassy, J.-P. Breittmayer and C. Aussel: Revaluation of the role of cholesterol in stabilizing rafts implicated in T cell receptor signaling. *Cellular Signalling*, 18(1), 105 (2006)
DOI: 10.1016/j.cellsig.2005.03.024
31. K. Gaus, M. Rodriguez, K. R. Ruberu, I. Gelissen, T. M. Sloane, L. Kritharides and W. Jessup: Domain-specific lipid distribution in macrophage plasma membranes. *J. Lipid Res.*, 46(7), 1526-1538 (2005)
DOI: 10.1194/jlr.M500103-JLR200
32. J. G. Kuhry, G. Duportail, C. Bronner and G. Laustriat: Plasma membrane fluidity measurements on whole living cells by fluorescence anisotropy of trimethylammoniumdiphenylhexatriene. *Biochim Biophys Acta*, 845, 60-7 (1985)
DOI: 10.1016/0167-4889(85)90055-2
33. L. Best, E. John and F. Jahning: Order and fluidity of lipid membranes as determined by fluorescence anisotropy decay. *European Biophysics Journal*, 15, 87-102 (1987)
DOI: 10.1007/bf00257502
34. B. R. Lentz: Use of fluorescent probes to monitor molecular order and motions within liposome bilayers. *Chemistry and Physics of Lipids*, 64, 99-116 (1993)
DOI: 10.1016/0009-3084(93)90060-G
35. A. Marczak: Fluorescence anisotropy of membrane fluidity probes in human erythrocytes incubated with anthracyclines and glutaraldehyde. *Bioelectrochemistry*, 74(2), 236 (2009)
DOI: 10.1016/j.bioelechem.2008.11.004
36. G. Gimpl, K. Burger and F. Fahrenholz: Cholesterol as modulator of receptor function. *Biochemistry*, 36, 10959-74 (1997)
DOI: 10.1021/bi963138w
37. J. Brejchová, J. Šádková, K. i. Dlouhá, L. Roubalová, P. Ostařov, M. Vořáhlíková, M. Hof and P. Svoboda: Fluorescence spectroscopy studies of HEK293 cells expressing DOR-Gi11 fusion protein; the effect of cholesterol depletion. *Biochim. et Biophys. Acta - Biomembranes*, 1808(12), 2819 (2011)
DOI: 10.1016/j.bbamem.2011.08.010
38. A. Gidwani, D. Holowka and B. Baird: Fluorescence anisotropy measurements of lipid order in plasma membranes and lipid rafts from rbl-2h3 mast cells. *Biochemistry*, 40, 12422-9 (2001)
DOI: 10.1021/bi010496c
39. S. Oncul, A. S. Klymchenko, O. A. Kucherak, A. P. Demchenko, S. Martin, M. Dontenwill, Y. Arntz, P. Didier, G. Duportail and Y. Maly: Liquid ordered phase in cell membranes evidenced by a hydration-sensitive probe: Effects of cholesterol depletion and apoptosis. *Biochim. et Biophys. Acta - Biomembranes*, 1798(7), 1436 (2010)
DOI: 10.1016/j.bbamem.2010.01.013
40. M. Korinek, V. Vyklicky, J. Borovska, K. Lichnerova, M. Kaniakova, B. Krausova, J. Krusek, A. Balik, T. Smejkalova, M. Horak and L. Vyklicky: Cholesterol modulates open probability and desensitization of NMDA receptors. *J. of Physiol*, 593(10), 2279 (2015)
DOI: 10.1113/jphysiol.2014.288209
41. M. Company, A. Iborra, J. Villaverde, P. Martínez and A. Morros: Membrane fluidity changes in goat sperm induced by cholesterol depletion using beta-cyclodextrin. *Biochim. Biophys. Acta - Biomembranes*, 1768(9), 2246 (2007)
DOI: 10.1016/j.bbamem.2007.06.011
42. T. J. Pucadyil and A. Chattopadhyay: Effect of cholesterol on lateral diffusion of fluorescent lipid probes in native hippocampal membranes. *Chemistry and Physics of Lipids*, 143(1-2), 11 (2006)
No DOI Found
43. C. Klein, T. Pillot, J. Chambaz and B. Drouet: Determination of plasma membrane fluidity with a fluorescent analogue of sphingomyelin by FRAP measurement using a standard confocal microscope. *Brain Research*

- Protocols*, 11(1), 46 (2003)
DOI: 10.1016/S1385-299X(03)00016-3
44. L. E. Organ and R. M. Raphael: Lipid lateral mobility in cochlear outer hair cells: regional differences and regulation by cholesterol. *J Assoc Res Otolaryngol.*, 10, 383-96 (2009)
DOI: 10.1007/s10162-009-0171-1
45. K. Gaus, E. Gratton, E. P. Kable, A. S. Jones, I. Gelissen, L. Kritharides and W. Jessup: Visualizing lipid structure and raft domains in living cells with two-photon microscopy. *Proc Natl Acad Sci U S A*, 100(26), 15554-9 (2003)
DOI: 10.1073/pnas.2534386100
46. K. Gaus, T. Zech and T. Harder: Visualizing membrane microdomains by Laurdan 2-photon microscopy. *Mol Membr Biol*, 23(1), 41-8 (2006)
DOI: 10.1080/09687860500466857
47. S. Y. Breusegem, N. Halaihel, M. Inoue, H. Zajicek, E. Lederer, N. P. Barry, V. Sorribas and M. Levi: Acute and chronic changes in cholesterol modulate Na-Pi cotransport activity in OK cells. *Am J Physiol Renal Physiol*, 289(1), F154-65 (2005)
DOI: 10.1152/ajprenal.00331.2004
48. S. Mukherjee and A. Chattopadhyay: Monitoring the organization and dynamics of bovine hippocampal membranes utilizing Laurdan generalized polarization. *Biochim. Biophys. Acta - Biomembranes*, 1714(1), 43 (2005)
DOI: 10.1016/j.bbamem.2005.06.010
49. T. P. Shentu, I. Titushkin, D. K. Singh, K. J. Gooch, P. V. Subbaiah, M. Cho and I. Levitan: oxLDL-induced decrease in lipid order of membrane domains is inversely correlated with endothelial stiffness and network formation. *Am J Physiol Cell Physiol*, 299(2), C218-229 (2010)
DOI: 10.1152/ajpcell.00383.2009
50. V. Borroni, C. J. Baier, T. Lang, I. Bonini, M. M. White, I. Garbus and F. J. Barrantes: Cholesterol depletion activates rapid internalization of submicron-sized acetylcholine receptor domains at the cell membrane. *Mol Membr Biol.*, 24, 1-15 (2007)
DOI: 10.1080/09687860600903387
51. J. Kwik, S. Boyle, D. Fooksman, L. Margolis, M. P. Sheetz and M. Edidin: Membrane cholesterol, lateral mobility, and the phosphatidylinositol 4,5-bisphosphate-dependent organization of cell actin. *Proc Natl Acad Sci U S A*, 100(24), 13964-9 (2003)
DOI: 10.1073/pnas.2336102100
52. K. M. S. O'Connell and M. M. Tamkun: Targeting of voltage-gated potassium channel isoforms to distinct cell surface microdomains. *J Cell Sci*, 118(10), 2155-2166 (2005)
DOI: 10.1242/jcs.02348
53. T. J. Pucadyil and A. Chattopadhyay: Cholesterol depletion induces dynamic confinement of the G-protein coupled serotonin1A receptor in the plasma membrane of living cells. *Biochim. Biophys. Acta - Biomembranes*, 1768(3), 655 (2007)
DOI: 10.1016/j.bbamem.2007.01.002
54. C. Baier, C. Gallegos, V. Levi and F. Barrantes: Cholesterol modulation of nicotinic acetylcholine receptor surface mobility. *European Biophysics Journal*, 39(2), 213-227 (2010)
DOI: 10.1007/s00249-009-0521-2
55. I. Levitan and K. J. Gooch: Lipid Rafts in Membrane-Cytoskeleton Interactions and Control of Cellular Biomechanics: Actions of oxLDL. *Antioxidants & Redox Signaling*, 9(9), 1519-1534 (2007)
DOI: 10.1089/ars.2007.1686
56. L. J. Pike and J. M. Miller: Cholesterol depletion delocalizes phosphatidylinositol bisphosphate and inhibits hormone-stimulated phosphatidylinositol turnover. *J. Biol. Chem.*, 273(35), 22298-22304 (1998)
DOI: 10.1074/jbc.273.35.22298
57. M. A. del Pozo, N. B. Alderson, W. B. Kiosses, H. H. Chiang, R. G. Anderson and M. A. Schwartz: Integrins regulate Rac targeting by internalization of membrane domains. *Science*, 303, 839-42 (2004)
DOI: 10.1126/science.1092571
58. T. Nebl, K. N. Pestonjamas, J. D. Leszyk, J. L. Crowley, S. W. Oh and E. J. Luna: Proteomic analysis of a detergent-resistant membrane skeleton from neutrophil plasma membranes. *J. Biol. Chem.*, 277(45), 43399-43409 (2002)
DOI: 10.1074/jbc.M205386200
59. M. Sun, N. Northup, F. Marga, F. J. Byfield, I. Levitan and G. Forgacs: Cellular cholesterol effects on membrane-cytoskeleton adhesion. *J Cell Sci.*, 120, 2223-31 (2007)

- DOI: 10.1242/jcs.001370
60. M. Sun, J. S. Graham, B. Hegedüs, F. Marga, Y. Zhang, G. Forgacs and M. Grandbois: Multiple membrane tethers probed by atomic force microscopy. *Biophys J*, 4320-9 (2005)
DOI: 10.1529/biophysj.104.058180
61. F. M. Hochmuth, J. Y. Shao, J. Dai and M. P. Sheetz: Deformation and flow of membrane into tethers extracted from neuronal growth cones. *Biophys. J.*, 70, 358–369 (1996)
DOI: 10.1016/S0006-3495(96)79577-2
62. Z. Hong, M. C. Staiculescu, P. Hampel, I. Levitan and F. Forgacs: How cholesterol regulates endothelial biomechanics?. *Frontiers in Vascular Biology*, 3, 426 (2012)
DOI: 10.3389/fphys.2012.00426
63. J. Pourati, A. Maniotis, D. Spiegel, J. L. Schaffer, J. P. Butler, J. J. Fredberg, D. E. Ingber, D. Stamenovic and N. Wang: Is cytoskeletal tension a major determinant of cell deformability in adherent endothelial cells? *Am J Physiol Cell Physiol*, 274(5), C1283-1289 (1998)
No DOI Found
64. F. Byfield, H. Aranda-Aspinoza, V. G. Romanenko, G. H. Rothblat and I. Levitan: Cholesterol depletion increases membrane stiffness of aortic endothelial cells. *Biophys J.*, 87, 3336-43 (2004)
DOI: 10.1529/biophysj.104.040634
65. Z. Hong, I. Ersoy, M. Sun, F. Bunyak, P. Hampel, Z. Hong, Z. Sun, Z. Li, I. Levitan, G. A. Meininger and K. Palaniappan: Influence of membrane cholesterol and substrate elasticity on endothelial cell spreading behavior. *J. of Biomedical Materials Research Part A*, 101A(7), 1994-2004 (2013)
DOI: 10.1002/jbm.a.34504
66. G. B. Kowalsky, D. Beam, M. J. Oh, F. Sachs, S. Z. Hua and I. Levitan: Cholesterol depletion facilitates recovery from hypotonic cell swelling in CHO cells. *Cell Physiol Biochem.*, 28, 1247-54 (2011)
DOI: 10.1159/000335856
67. F. J. Byfield, B. D. Hoffman, V. G. Romanenko, Y. Fang, J. C. Crocker and I. Levitan: Evidence for the role of cell stiffness in modulation of volume-regulated anion channels. *Acta Physiologica*, 187(1-2), 285-294 (2006)
DOI: 10.1111/j.1748-1716.2006.01555.x
68. T. G. Mason and D. A. Weitz: Optical Measurements of Frequency-Dependent Linear Viscoelastic Moduli of Complex Fluids. *Physical Review Letters*, 74(7), 1250-1253 (1995)
DOI: 10.1103/PhysRevLett.74.1250
69. J. C. Crocker and D. G. Grier: Methods of digital video microscopy for colloidal studies. *J. of Colloid and Interface Science*, 179(1), 298-310 (1996)
DOI: 10.1006/jcis.1996.0217
70. A. W. C. Lau, B. D. Hoffman, A. Davies, J. C. Crocker and T. C. Lubensky: Microrheology, stress fluctuations, and active behavior of living cells. *Physical Review Letters*, 91(19), 198101-(1-4) (2003)
No DOI Found
71. J. C. Crocker, M. T. Valentine, E. R. Weeks, T. Gisler, P. D. Kaplan, A. G. Yodh and D. A. Weitz: Two-point microrheology of inhomogeneous soft materials. *Physical Review Letters*, 85(4), 888-891 (2000)
DOI: 10.1103/PhysRevLett.85.888
72. F. J. Byfield, S. Tikku, G. H. Rothblat, K. J. Gooch and I. Levitan: OxLDL increases endothelial stiffness, force generation and network formation. *J. Lipid Res.*, 47, 715-723 (2006)
DOI: 10.1194/jlr.M500439-JLR200
73. A. L. Sieminski, R. P. Hebbel and K. J. Gooch: The relative magnitudes of endothelial force generation and matrix stiffness modulate capillary morphogenesis *in vitro*. *Exp Cell Res.*, 297, 574-84 (2004)
DOI: 10.1016/j.yexcr.2004.03.035
74. M. Dembo and Y. L. Wang: Stresses at the cell-to-substrate interface during locomotion of fibroblasts. *Biophys J.*, 76, 2307-16 (1999)
DOI: 10.1016/S0006-3495(99)77386-8
75. L. L. Norman, R. J. Oetama, M. Dembo, F. Byfield, D. A. Hammer, I. Levitan and H. Aranda-Espinoza: Modification of Cellular Cholesterol Content Affects Traction Force, Adhesion and Cell Spreading. *Cell Mol Bioeng.*, 3, 151-162 (2010)
DOI: 10.1007/s12195-010-0119-x
76. K. M. Stroka, I. Levitan and H. Aranda-Espinoza: OxLDL and substrate stiffness promote neutrophil transmigration by enhanced endothelial cell contractility

- and ICAM-1. *J. of Biomechanics*, 45(10), 1828 (2012)
DOI: 10.1016/j.jbiomech.2012.04.011
77. G. B. Kowalsky, F. J. Byfield and I. Levitan: oxLDL facilitates flow-induced realignment of aortic endothelial cells. *Am J Physiol Cell Physiol*, 295(2), C332-340 (2008)
DOI: 10.1152/ajpcell.00335.2007
78. T. P. Shentu, D. K. Singh, M.-J. Oh, S. Sun, L. Sadaat, A. Makino, T. Mazzone, P. V. Subbaiah, M. Cho and I. Levitan: The role of oxysterols in control of endothelial stiffness. *J. of Lipid Res.*, 53., 1348-58 (2012)
DOI: 10.1194/jlr.M027102
79. S. Yla-Herttuala, W. Palinski, M. E. Rosenfeld, S. Parthasarathy, T. E. Carew, S. Butler, J. L. Witztum and D. Steinberg: Evidence for the presence of oxidatively modified low density lipoprotein in atherosclerotic lesions of rabbit and man. *J Clin Invest.*, 84, 1086-95 (1989)
DOI: 10.1172/JCI114271
80. L. J. van Tits, T. M. van Himbergen, H. L. Lemmers, J. de Graaf and A. F. Stalenhoef: Proportion of oxidized LDL relative to plasma apolipoprotein b does not change during statin therapy in patients with heterozygous familial hypercholesterolemia. *Atherosclerosis*, 185, 307-12 (2005)
DOI: 10.1016/j.atherosclerosis.2005.06.006
81. P. Holvoet, A. Mertens, P. Verhamme, K. Bogaerts, G. Beyens, R. Verhaeghe, D. Collen, E. Muls and F. Van de Werf: Circulating Oxidized LDL Is a Useful Marker for Identifying Patients With Coronary Artery Disease. *Arterioscler Thromb Vasc Biol*, 21(5), 844-848 (2001)
DOI: 10.1161/01.ATV.21.5.844
82. I. Levitan, S. Volkov and P. V. Subbaiah: Oxidized LDL: Diversity, patterns of recognition and pathophysiology. *Antioxidants & Redox Signaling*, 13, 39-75 (2010).
DOI: 10.1089/ars.2009.2733
83. A. Blair, P. W. Shaul, I. S. Yuhanna, P. A. Conrad and E. J. Smart: Oxidized low-density lipoprotein displaces endothelial Nitric-oxide synthase from plasmalemmal caveolae and impairs eNOS activation. *The J. of Biol. Chem.*, 274, 32512-32519 (1999)
DOI: 10.1074/jbc.274.45.32512
84. X. Xu and E. London: The effect of sterol structure on membrane lipid domains reveals how cholesterol can induce lipid domain formation. *Biochemistry*, 39, 843-849 (2000)
DOI: 10.1021/bi992543v
85. H. Oberleithner, C. Riethmuller, H. Schillers, G. A. MacGregor, H. E. de Wardener and M. Hausberg: Plasma sodium stiffens vascular endothelium and reduces nitric oxide release. *Proc Natl Acad Sci U S A*, 104(41), 16281-6 (2007)
DOI: 10.1073/pnas.0707791104
86. F. J. Byfield, R. K. Reen, T.-P. Shentu, I. Levitan and K. J. Gooch: Endothelial actin and cell stiffness is modulated by substrate stiffness in 2D and 3D. *J. of Biomechanics*, 42(8), 1114 (2009)
DOI: 10.1016/j.jbiomech.2009.02.012
87. M. S. Gundavaram, T. P. Shentu, G. B. Kowalsky, S. Volkov, D. E. Schraufnagel and I. Levitan: Oxidized low-density lipoprotein alters the effect of matrix stiffness on the formation of endothelial networks and capillary lumens. *Pulmonary Circulation*, 3, 622-31 (2013)
DOI: 10.1086/674309
88. N. L. Sehgel, Y. Zhu, Z. Sun, J. P. Trzeciakowski, Z. Hong, W. C. Hunter, D. E. Vatner, G. A. Meininger and S. F. Vatner: Increased vascular smooth muscle cell stiffness: a novel mechanism for aortic stiffness in hypertension. *Am. J. of Physiol - Heart and Circulatory Physiology*, 305(9), H1281-H1287 (2013).
DOI: 10.1152/ajpheart.00232.2013
89. N. L. Sehgel, Z. Sun, Z. Hong, W. C. Hunter, M. A. Hill, D. E. Vatner, S. F. Vatner and G. A. Meininger: Augmented Vascular Smooth Muscle Cell Stiffness and Adhesion When Hypertension Is Superimposed on Aging. *Hypertension*, 65(2), 370-377 (2015)
DOI: 10.1161/HYPERTENSIONAHA.114.04456

Key Words: Cell stiffness, Cell Biomechanics, Oxidized Lipids, Review

Send correspondence to: Irena Levitan, Departments of Medicine, Pharmacology and Bioengineering, University of Illinois at Chicago, 909 S. Wolcott Ave., Bldg. COMRB RM3097, Chicago, IL 60612, Tel: -312-996-0441, Fax: 312-996-7193, E-mail: levitan@uic.edu